

A note on price adjustment with menu cost for multi-product firms*

Fernando Alvarez

University of Chicago

Francesco Lippi

University of Sassari, EIEF

November 15, 2010

PRELIMINARY DRAFT

Abstract

We study the stylized problem of a multi-product firm that can revise prices only after paying a fixed “menu” cost. The key assumption, introduced by [Lach and Tsiddon \(1996, 2007\)](#) and [Midrigan \(2007, 2009\)](#), is that once the menu cost is paid the firm can adjust the price of *all* its products. The firm’s problem is to minimize the deviations of the profits incurred relative to the flexible price case, i.e. the case with no menu cost. We completely characterize the solution of a simple symmetric problem in terms of the structural parameters: the variability of the flexible prices, the curvature of the profit function, the size of the menu cost, and the number of products sold by the firm. We also provide analytical expressions for the frequency of adjustment, the hazard rate of price adjustments, and the distribution of price changes in terms of the structural parameters.

JEL Classification Numbers: E5

Key Words: menu cost, economies of scope in price changes, optimal control subject to fixed cost in multiple dimensions

*We thank

1 Introduction

Several papers document how price setting behavior, as summarized by the size-distribution and by the timing of price changes, varies systematically with the number of products sold.¹ The recent empirical work of [Bhattarai and Schoenle \(2010\)](#) documents that firms selling more goods display a higher frequency of price adjustment. [Lach and Tsiddon \(2007\)](#) show that price changes are synchronized within stores, but staggered across stores.

Despite the rich and growing evidence on this phenomenon, there is scant theoretical work on this problem. [Midrigan \(2007, 2009\)](#) begun to study this problem by explicitly writing down and solving numerically a model where a firm is selling 2 goods, subject to a common menu cost.² Compared to the classic case of one good, his model generates a distribution of price changes with “small” price adjustments. But some important questions remain to be answered: what forces shape the optimal pricing decisions as the number of goods n sold by the firm changes? going beyond the $n = 2$ case is important, as the number of goods sold by the retail stores, where much of the micro data are measured, is much larger.³

This paper provides a simple model to study how the price setting decision depends on n , the number of goods sold by a firm. The model allows us to answer some questions that are hard to tackle without a formal frame. Examples of these questions are: Is the higher frequency of price adjustment the mechanical consequence of a fixed menu ψ cost split over a larger number of goods, so that it is as if each good pays a smaller menu cost ψ/n as n increases? in other words, how does the frequency of price adjustment behaves as a function of n if the menu cost paid by the firm increased linearly with the number of goods: $n \psi$? Is the synchronization of price adjustment arising from the fact that a single fixed cost applies

¹ An incomplete list of contributions documenting this fact includes [Lach and Tsiddon \(1992\)](#), [Lach and Tsiddon \(2007\)](#), [Baudry et al. \(2007\)](#), [Dhyne and Konieczny \(2007\)](#), [Dutta et al. \(1999\)](#), [Midrigan \(2007, 2009\)](#), [Cavallo \(2009\)](#), [Bhattarai and Schoenle \(2010\)](#) and [Neiman \(2010\)](#).

² Importantly, he also put this model into a general equilibrium framework, and analyzed the effect of a monetary shock. [Bhattarai and Schoenle \(2010\)](#) also solve numerically the problem for a firm selling three goods.

³[Bhattarai and Schoenle \(2010\)](#) reports that the US the median firm sampled by the BLS sells between 3 to 5 goods.

to a bundle of goods, or is there a systematic association between the number of products and the volatility σ of the shock that hit each product? One purpose of this note is to provide some guidance for the empirical examination of these concerns by analyzing the identification of the different factors, such as the n, σ^2, ψ , and B in a tractable set up. Another purpose of the note is to advance on the understanding of the impact effect of an aggregate monetary shock in this set-up. This question that has been tackled numerically by [Midrigan \(2007\)](#) for the case of $n = 2$, but as we explain below we provide tools that help to analyze in the case of $n > 2$.

We study a stylized version of the problem of a multi-product firm that can revise prices only after paying a fixed cost. The key assumption, introduced by [Lach and Tsiddon \(1996, 2007\)](#) and [Midrigan \(2007, 2009\)](#), is that once the fixed menu cost is paid the firm can adjust the price of *all* its products. The problem is set up as to minimize the deviations of the profits incurred relative to the flexible price case, i.e. the case with no menu cost. We assume that the static profit maximizing price for each of the n products, which coincide with the price that will be charged without menu cost, follows n independent random walks without drift and with volatility σ per unit of time. We refer to the vector of the difference between the frictionless prices and the actual prices charged as the vector of the price gaps. The period return function is assumed to be proportional to the sum of the squares of the price gaps. The proportionality constant B measures the second order per period losses associated with charging a price different from the optimum, i.e. it is a measure of the curvature of the profit function.⁴ We assume that if a fixed cost ψ is paid the firm can simultaneously change all the prices. The firm minimizes the expected discounted cost, which include the stream of lost profit from charging prices different from the frictionless as well as the fixed cost at the time of adjustments. We completely characterize the solution of a simple symmetric problem in terms of the structural parameters: the variability of the flexible prices σ , the curvature of the profit function B , the size of the menu cost ψ , the discount rate r , and the number of

⁴The first order losses are zero, since the maximum per period profits are obtained at the frictionless price.

products n . We also provide analytical expressions for the invariant distribution of the price gaps, the frequency of adjustment, the hazard rate of price adjustments, and the marginal distribution of price changes in terms of the fundamental parameters.

The solution of the firms' problem involves finding the set over which prices are adjusted, and the set where they are not, i.e. the inaction set. Due to the lack of drift, when prices are adjusted they are set equal to the frictionless prices, i.e. the price gaps are set to zero in all dimensions. We show that the optimal decision is to control the price gap as to remain in the interior of the n -dimensional ball centered at the origin. The size of this ball, whose square radius we denote by \bar{R} , is chosen optimally. We solve for the value function and completely characterize the size of the inaction set \bar{R} as a function of the parameters of the problem. As we let $r \downarrow 0$ the ratio \bar{R}/σ^2 can be written as an increasing function of two arguments: $\sigma^2 B/\psi$ and n . We also obtain a very accurate approximation for small cost ψ , where we show that \bar{R} takes the form of a square root function, $\bar{R} \approx [2(n+2)\sigma^2 B/\psi]^{1/2}$.

To our knowledge this is the first fixed cost adjustment problem in n -dimensions whose solution is analytically characterized. We believe that this is because of the difficulty of finding a tractable boundary condition and a candidate solution that is smooth enough on the boundary of the inaction region. [Baccarin \(2009\)](#) gives a recent statement of the general problem, and a characterization of the viscosity solution. In our case we can reduce the dimension of the problem to one, by keeping track of R , the square of the radius of the vector of the price gaps. This reduction is possible because of the quadratic nature of the objective function, and the lack of drift of the uncontrolled price gap. Thus, we trade off high dimension for a non-linearity on the evolution of the system.

As mentioned, [Midrigan \(2007, 2009\)](#) analyzes the effect of monetary policy shock in a general equilibrium model with $n = 2$. He shows that it differs from the one with $n = 1$ because, among other things, the mass of firms that are close to the inaction region just before the shock and find it optimal to adjust right after the shock is smaller when $n = 2$. The reason is that, for $n = 2$, some price adjustments are small because not all individual

prices being adjusted are close to the inaction boundary. In the language of Golosov and Lucas (2007), there is a smaller selection effect among the firms that adjust in the $n = 2$ case, compared to the $n = 1$ case. Indeed Caballero and Engel (2007) argue that the cross section distribution of the “desired adjustments”, or price gaps in our set up, is one of the key ingredients to understand the aggregate effect in a model with sS policies. Motivated by these findings, we compute the density of the invariant distribution of R , the sum of the square of the price gaps. This density is downward sloping, and, except for its scale, it depends *exclusively* on the number of products n . For larger values of n this distribution puts relatively more mass on the points close to the boundary of the sphere, i.e. there are more firms close to the point where they want to adjust. As $n \rightarrow \infty$, the distribution is uniform. [WE NEED TO FINISH THIS PARAGRAPH...]

We characterize the implications for the timing of price changes given \bar{R} , σ^2 and n . We show that the expected number of price adjustments per unit of time is given by $n\sigma^2/\bar{R}$, which together with our result for \bar{R} gives a complete characterization of the frequency of price adjustments. This characterization can be used to disentangle the effects on the frequency of adjustments while comparing firms with different number of products, since it points out to all its determinants. Moreover, when used together with other information described below, it can be used to identify the parameters of the model and test its implications.

We solve in closed form for the hazard rate of the price changes as a function of the time elapsed since the last change. The shape of this function, except for its scale, depends *exclusively* on the number of products n . The scale of the function is completely determined by the expected number of adjustment per unit of time, which we have already solve for. For a given n , the hazard rates are increasing in duration, have an elongated S shape, with a finite asymptote. Comparing across different values of n , while keeping the expected number of adjustment constant, we show that the asymptote is increasing in n . As we let n increase without bound, the asymptote diverges to $+\infty$ and the hazard rate function converges to the one with deterministic adjustments, i.e. towards one with an inverted L shape.

Finally we characterize the shape of the distribution of price changes. While price changes occur simultaneously for n products, we characterized the marginal distribution of prices, because this is the object that is usually computed in actual data sets. We give a close form expression for the density of the marginal distribution of price changes as a function of \bar{R} and n . Thus, once the size of the changes is control for, the shape of the price change distribution is *exclusively* a function of the number of products. We obtain then several statistics that have been computed in the data, such as the coefficient of variation of the absolute value of price changes, or the excess kurtosis, as purely functions of the n . Indeed the shape of the distribution of price changes is as follows: for $n = 2$ it is bimodal, with modes at the absolute value of $\sqrt{\bar{R}}$, for $n = 3$ is uniform, for $n = 4$ peaks at zero and it is concave, and for larger n it is bell shape. Indeed, as $n \rightarrow \infty$, once normalized, the distribution converged to a standard normal.

2 A stylized multiproduct menu cost model

Let n be the number of goods produced by the firm. Each price p_i evolves according to a random walk without drift, so that $dp_i = \sigma dW_i$ where dW_i is a standard Brownian Motion. The n Brownian Motions (BM henceforth) are independent, so $\mathbb{E}[W_i(t)W_j(t')] = 0$ for all $t, t' \geq 0$ and $i, j = 1, \dots, n$. The problem is:

$$V(p) = \min_{\{\tau_j, u(\tau_j)\}_{j=1}^{\infty}} \mathbb{E} \left[\sum_{j=1}^{\infty} e^{-r\tau_j} \psi + \int_0^{\infty} e^{-rt} B \left(\sum_{i=1}^n p_i^2(t) \right) dt \middle| p(0) = p \right] \quad (1)$$

where

$$p_i(t) = \sigma W_i(t) + \sum_{j:\tau_j < t} \Delta p_i(\tau_j) \text{ for all } t \geq 0 \text{ and } i = 1, 2, \dots, n, \quad (2)$$

and $p(0) = p$.

So that τ_j are the (stopping) times at which control is exercised. At these times, after paying the cost ψ , the state can be changed to any value in \mathbb{R}^n . We denote the vector of price

changes as $\Delta p(\tau_j) \in \mathbb{R}^n$. This is a standard adjustment cost problem subject to a fixed cost, with the exception that after paying the adjustment cost ψ the decision maker can adjust the state in the n dimension.

At an abstract level [equation \(1\)](#) and [equation \(2\)](#) can be used to solve a symmetric quadratic loss tracking problem in n dimensions, subject to a fixed adjustment cost. To map it into a tracking problem, let the state of the system be two n dimensional vectors $\hat{p}(t)$, and $p^*(t)$. The interpretation of $\hat{p}(t)$ is the location of the system, and $p^*(t)$ a bliss point, the location that the decision maker is tracking. The instantaneous cost of the decision maker is proportional to the distance between the location of the system and the bliss point, $B\|p^*(t) - \hat{p}(t)\|^2$, where $B > 0$. Each component of the bliss points evolve as an independent random walk without drift, with variance σ^2 per unit of time. If the decision maker pays a fixed cost ψ/n she can change the location of the system anywhere that she desires. For the purpose of finding the times at which the decision maker chooses to change the state, and to find the value of the changes of the state, we can simplify the problem and consider the distance between the location of the state and the bliss point the of the system, and simple let the state be $p(t) = \hat{p}(t) - p^*(t)$. We have written [equation \(1\)](#) and [equation \(2\)](#) using this “gap” notation.

For future reference we note if B and ψ are multiplied by a constant $\lambda > 0$ the value function is scaled by λ with no change on the decisions. This explains why all the decision are functions of B/ψ . We will use this property to interpret different assumptions about how these parameter vary firms with different number of products n .

We describe now an economic interpretation of the problem, which can be summarized to say that the firms “tracks” the prices that will maximize instantaneous profits from the n products. Consider a system of n independent demands, with constant elasticity η for each product, and a time varying constant marginal cost $C_i(t)$. In the context of the price setting models, our model is a stylized version of the problem introduced by [Midrigan \(2007, 2009\)](#) where the elasticity of substitution between the goods produced *within* the firm is the same as

the one of the bundle of goods produced *across* firms. The instantaneous profit maximizing price is proportional to the marginal cost, or in logs $p_i^*(t) = \log c_i(t) + \log((\eta - 1)/\eta)$. In this case we assume that the log of the marginal cost evolve as a random walk with drift so that $p_i^*(t)$ inherits this property. We can interpret the period cost as a second order expansion of the (log) of the profit function with respect to the vector of the log of prices, around the log of the profit maximized price vector. The first order term are zero because we are expanding around $p^*(t)$. The fact that it is an expansion of the *log* of the profit is equivalent to measure the profits relative to the value of the maximized profit for the n goods. There are no second order cross terms due to the separability of the demand. Thus we can write the problem in terms of the gap between the actual price and the profit maximizing price: $p(t) = \hat{p}(t) - p^*(t)$. The constant B is given by $B = (1/2)\eta(\eta - 1)/n$, where the term $1/2$ is due to a second order expansion, the terms with η are due to the fact that the curvature of the profits depend on the elasticity of demand, and the term $(1/n)$ is the the share of profits from each product relative to the profits across the n goods. In this interpretation the value of the fixed cost is measured relative to the profit of the n goods, thus it costs ψ/n in units of the numeraire good. Since all that matter for the decision is the ratio of B to ψ this normalization only scales the units of the vale function. In [Appendix C](#) we give a derivation of the second order expansion as well as a discussion and interpretation of the parameters B and ψ . In particular the interpretation of how to scale B and ψ for firms with different of products.

Below we consider two cases for scaling of the cost of adjustment with respect to the number of goods. In the first case, which we refer to as constant returns to scale (CRTS) technology for adjustment cost, when we compare firms with different values of n , the adjustment cost scales linearly with it, so that $\psi = n \psi_1$. In this case a firm with twice as many products pays twice as much in terms of the numeraire good to adjust all the prices simultaneously. We refer to the case of a *constant fixed cost*, if $\psi = \psi_1$, so that a firm with twice as many products pays the same cost in terms of numeraire to adjust the twice as many prices. We think that these two extreme simple cases bracket all of the interesting setups.

We note the following basic properties of the value function and the optimal policy.

1. Given the symmetry of the BM and of the objective function around zero, and the independence of the BM's, one can use reflection around zero to show that the value function only depends on the absolute values of p_i , i.e. $V(p) = V(|p_1|, |p_2|, \dots, |p_n|)$ for all $p \in \mathbb{R}^n$.
2. Due to the the symmetry of the problem, and the shape of the period return function, we have that after an adjustment the state is reset at the origin, i.e. $p(\tau_j^+) = 0$, or $\Delta p(\tau_j) = -p(\tau_j^-)$.
3. The state space \mathbb{R}^n can be divided in two regions, an inaction region $\mathcal{I} \subset \mathbb{R}^n$ and control region $\mathcal{C} \subset \mathbb{R}^n$. We use $\text{Int}(\mathcal{I})$ for the interior of the range of inaction, and $\partial\mathcal{I}$ for its boundary. We have that $\mathcal{C} \cap \text{Int}(\mathcal{I}) = \emptyset$, that inaction is strictly preferred in $\text{Int}(\mathcal{I})$, that control is strictly preferred in $\text{Int}(\mathcal{C})$, and that in $\partial\mathcal{I}$ the agent is indifferent between control and inaction.

We write down the conditions for the solution of the problem, provided that a value function is smooth enough (i.e. we look for a solution of the “strong” formulation of the problem). In the range of inaction the cost for the firm is given by the following Bellman equation:

$$r V(p_1, p_2, \dots, p_n) = B \sum_{i=1}^n p_i^2 + \frac{\sigma^2}{2} \sum_{i=1}^n V_{ii}(p_1, p_2, \dots, p_n) \quad (3)$$

for all $p \in \mathcal{I}$. In the control region we have:

$$V(p_1, p_2, \dots, p_n) = V(0) + \psi \quad (4)$$

for all $p \in \mathcal{C}$. The optimality of returning to the origin implies that,

$$V_i(p_1, p_2, \dots, p_n) = 0 \text{ for all } i = 1, 2, \dots, n \quad (5)$$

and that $V(p)$ is convex in a neighborhood of the origin. Finally, differentiability in the boundary of the inaction

$$V_i(p_1, p_2, \dots, p_n) = 0 \text{ for } i = 1, 2, \dots, n \text{ and for all } p \in \partial\mathcal{I}, \quad (6)$$

We refer to this condition as smooth pasting.

3 Characterization of the solution

Before presenting the solution of this problem we change the state space, which we summarize using a single variable. Let

$$R = \sum_{i=1}^n p_i^2 \quad (7)$$

measure the deviation of prices from their optimal value across the n goods. We consider policies summarized by a single number \bar{R} . In this class of policies the firm controls the state so that if $R < \bar{R}$, there is inaction. The first time that R reaches \bar{R} , all prices are adjusted to the origin, so that $R = 0$. We will find the optimal policy in this class. Then we will show that the optimal policy of the original problem is of this form.

The variable R measures the square of the ray of a sphere centered on the origin. Since each of the prices follows identical independent standard BM in the inaction region, then R follows a simple diffusion in the inaction. Using Ito's Lemma on [equation \(7\)](#) the evolution of R is

$$dR = n\sigma^2 dt + 2\sigma \sum_{i=1}^n p_i(t) dW_i$$

This implies that the quadratic variation of R is:

$$\mathbb{E}(dR)^2 = 4\sigma^2 \left(\sum_{i=1}^n p_i^2(t) \right) dt$$

Thus we can define a stochastic differential equation for R with a new standard BM $\{W(t)\}$

that solves:

$$dR = n\sigma^2 dt + 2\sigma\sqrt{\bar{R}} dW \text{ for } R \in [0, \bar{R}]. \quad (8)$$

We note that for the unregulated process, i.e. when $\bar{R} = \infty$, if $R(0) > 0$ then $R(t) > 0$ for $t > 0$ with probability one provided that $n \geq 2$, see [Karatzas and Shreve \(1991\)](#) Proposition 3.22.⁵

Note that the drift and diffusion terms in [equation \(8\)](#) are only functions of R . We also note that instantaneous return is a function of R , so we can write the following

$$v(R) = \min_{\bar{R}} \mathbb{E} \left[\sum_{j=1}^{\infty} e^{-r\tau_j} \psi + \int_0^{\infty} e^{-rt} B R(t) dt \mid R(0) = R \right] \quad (9)$$

subject to [equation \(8\)](#) when $R \in [0, \bar{R}]$, where τ_j are the first time that $R(t)$ hits \bar{R} . The function v solves:

$$r v(R) = BR + n\sigma^2 v'(R) + 2\sigma^2 R v''(R), \quad \text{for } R \in (0, \bar{R}) . \quad (10)$$

Since policy calls for adjustment at values higher than \bar{R} we have:

$$v(R) = v(0) + \psi, \quad \text{for all } R \geq \bar{R} . \quad (11)$$

If v is differentiable at \bar{R} we can write the two boundary conditions:

$$v(\bar{R}) = v(0) + \psi \quad \text{and} \quad v'(\bar{R}) = 0 . \quad (12)$$

These conditions are typically referred to as value matching and smooth pasting. For $R = 0$

⁵This result was obtained for Bessel processes, which are the square root of $R(t)$. Additionally, [Karatzas and Shreve \(1991\)](#) have shown in Problem 3.23 and 3.24 that for if $R(0) > 0$, then for $n = 2$ the unregulated process can become arbitrarily close to zero but for $n \geq 3$ almost every path remains bounded away from zero. Furthermore, for the regulated process the classification for the boundaries of a diffusion gives that for $n \geq 2$ the point $R = 0$ is an entrance boundary, as verified in [Karlin and Taylor \(1999\)](#) Example 6, Chapter 12.6.

to be the optimal return point, it must be a global minimum, and thus we require that:

$$v'(0) \geq 0 . \tag{13}$$

Note the weak inequality, since R is non-negative.

The next proposition finds an analytical solution for v in the range of inaction.

PROPOSITION 1. Let $\sigma > 0$. The ODE equation (10) has the solution:

$$v(R) = \sum_{i=0}^{\infty} \beta_i R^i , \quad \text{for } R \in [0, \bar{R}] . \tag{14}$$

where the coefficients $\{\beta_i\}$ solve:

$$\beta_0 = \frac{n\sigma^2}{r}\beta_1 , \quad \beta_2 = \frac{r\beta_1 - B}{2\sigma^2(n+2)} , \quad \beta_{i+1} = \frac{r}{(i+1)\sigma^2(n+2i)}\beta_i , \quad \text{for } i \geq 2 . \tag{15}$$

for any β_0 .

The proof follows by replacing the function given in equation (14) into the ODE (10) and matching the coefficients for the powers of R^i . The next proposition shows that there exist a unique solution of the ODE (10) satisfying the relevant boundary conditions. It partially characterizes the solution of the threshold \bar{R} .

PROPOSITION 2. Assume $r > 0, \sigma > 0, n \geq 1$. There exists a unique solution of the ODE (10) satisfying the boundary conditions equation (12) for which $v(\cdot)$ is minimized at $R = 0$. The unique value of $\bar{R}(\psi)$ is strictly increasing in ψ , with $\bar{R}(0) = 0$ and $\bar{R} \rightarrow \infty$ as $\psi \rightarrow \infty$. Moreover, as $r \downarrow 0$, the optimal threshold satisfies $\bar{R} = \sigma^2 q\left(\frac{\psi}{B\sigma^2}, n\right)$ for some strictly increasing function $q(\cdot, n)$.

We note that the solution for \bar{R} is only a function of the ratio ψ/B is apparent from the definition of the sequence problem. That it is strictly increasing in the ratio of the fixed cost to the benefit of adjustment ψ/B is quite intuitive. The last property implies that if \bar{R} is

the solution of the problem for parameters $\sigma, \psi/B$ and n , then $\bar{R}' = \bar{R} \lambda$ is the solution for parameters $(\psi/B)' = \lambda, n' = n$ and $\sigma' = \sqrt{\lambda} \sigma$ for any $\lambda > 0$.

While the last proposition gives a partial characterization of the optimal thresholds, it does not analyze how it depends on n . To do so, we give a simple expression for an approximate solution of \bar{R} . The formula gives an approximate solution for small values of ψ and r . Mechanically we take value matching and smooth pasting conditions derived above and we set $\beta_i = 0$ for $i \geq 3$, i.e. we assume that $v(\cdot)$ is quadratic. Taking r to zero gives:

$$\bar{R} = \sqrt{\frac{\psi \sigma^2 2 (n + 2)}{B}}. \quad (16)$$

Notice that this approximation satisfied the properties of the general solution of [Proposition 2](#). The effect on this formula of $\psi\sigma^2/B$ is exactly the same as in the case of one product. Indeed the quartic root implied when $n = 1$ was obtained by [Dixit \(1991\)](#) for the model with $n = 1$. We found that the quadratic approximation to $v(\cdot)$, which amounts to a quartic approximation to $V(\cdot)$, gives very accurate values for \bar{R} . Alternatively, the rest of the β_i coefficients become very small for $i \geq 3$ for the parameters that we are interested.

The next proposition verifies that the restricted policy that we use to characterize the solution is indeed optimal. Indeed the decision rule is very similar to the one obtained by [Baccarin \(2009\)](#). He studies a general class of problems with fixed and variable adjustment cost and a non-linear cost function, but illustrates his results with the computation of several examples, among them one very similar to ours.

PROPOSITION 3. Let v be the solution of the restricted problem [equation \(9\)](#) and [equation \(8\)](#). Let $V(p) = v(\sum_{i=1}^n p_i^2)$. This is the solution of the problem described [equation \(1\)](#) and [equation \(2\)](#).

For completeness we note that in the case of $n = 1$ and the case of perfectly correlated target prices. In the case of one product, i.e. $n = 1$, the solution to V is easily seen to be

the sum of a quadratic and an two exponential, as in

$$V(p) = \frac{B}{r}p^2 + \beta (\exp(\lambda p) + \exp(-\lambda p)) + B \left(\frac{\sigma}{r}\right)^2$$

where $\lambda = \sqrt{2r}/\sigma$ and the constant β is chosen to match enforce smooth pasting and value matching. Moreover, it is easy to see that in that case the $v(R) = V(\sqrt{R})$ solves the ODE in (10) and its boundary conditions. We note that the solution for the $n = 1$ case and the expression for the approximation for \bar{R} are the same as the ones derived in Dixit (1991), which we explore in the price setting context in Alvarez, Lippi, and Paciello (2009). In the case of n perfectly correlated target prices the problem has, after the first adjustment, a single state variable. In this case, in terms of the threshold policy and value function, the problem is identical to the one with only one price. The static return is thus $nBp(t)^2$ where $p(t)$ is, when uncontrolled, a one dimensional brownian motion. The only difference with the problem with only one price is that the value of B is multiplied by n , or more importantly, the ratio B/ψ is proportional to n . This is quite natural, since the adjustment has the same effectiveness for all products, and hence it is as if it were cheaper. Note that, in the case of the CRTS assumption, the value of the adjustment threshold, and hence the frequency of adjustment, is independent of n . Instead, in terms of the implication for price changes, the problem with perfectly correlated shocks is quite differently, since there are *no* small price changes. When adjustment takes place, all products have the same price gap. We return to this simple case later on to speculate on the case of positive, but less than one correlation between the innovations.

4 Implications for timing and size of price changes

In this section we explore the implications for the frequency and distribution of price changes.

We let the expected time for $R(t)$ to hit the barrier \bar{R} starting at R by the function $\mathcal{T}(R)$.

This function satisfies:

$$0 = 1 + n \sigma^2 \mathcal{T}'(R) + 2 R \sigma^2 \mathcal{T}''(R) \text{ for } R \in (0, \bar{R}) \text{ and } \mathcal{T}(\bar{R}) = 0 ,$$

where the first condition gives the law of motion inside the range of inaction and the second one imposes the terminal condition on the boundary of the range of inaction. The unique solution of this ODE that satisfies the relevant boundary condition is:

$$\mathcal{T}(R) = \frac{\bar{R} - R}{n \sigma^2} \text{ for } R \in [0, \bar{R}] . \quad (17)$$

We use $\mathcal{T}(0)$ as the expected time between successive price adjustments, and thus the average number of adjustment, denoted by N_a is given by $\frac{1}{\mathcal{T}(0)}$. We collect this result in the following proposition:

PROPOSITION 4. Let N_a be the expected number of price changes for a multi-product firm with n goods. It is given by

$$N_a = \frac{n\sigma^2}{\bar{R}} \cong \sqrt{\frac{B \sigma^2}{2 \psi} \frac{n^2}{(n+2)}} . \quad (18)$$

In the last expression of [equation \(18\)](#) we use the approximation of \bar{R} for small ψ and r . It is interesting that this expression extends a well known expression for the case of $n = 1$, simply by adjusting the value of the variance from σ^2 to $n\sigma^2$. The number of products n affects N_a through two opposing forces. One is that with more products, the variance of the deviations of the price gaps increases, and thus a given value of \bar{R} is hit sooner in expected value, which we refer to as the direct effect. On the other hand, with with more products, the optimal value of \bar{R} is higher. Expression [equation \(18\)](#) shows that, as often happens in these models, the direct effect dominates, and the frequency of adjustment increases with n .

We use this expression to study how the bundling of menu costs, i.e. the fact that a single menu cost relates to several products, affects the frequency of adjustment of individual

prices. This is interesting because recent evidence in [Bhattarai and Schoenle \(2010\)](#) shows that the the frequency of price adjustment appears higher for firms that sell a larger number of goods.⁶ They find that the average frequency of price adjustment increases, from about 20% to about 30% per month, moving from the first to the fourth bin.

This pattern is qualitatively consistent with the formula in [equation \(18\)](#), which shows that N_a is increasing in n . Notice however that in this comparison we are keeping ψ constant, so that as n increases the menu cost per good is decreasing. One may wonder whether the increased activity by the firms follows from the fact that the menu cost is smaller (per good) or because of the bundling of the goods prices. To separate the effects of the economies of scale in the menu cost from the bundling of the goods, consider the case where the cost ψ grows linearly with the number of goods n , i.e. : $\psi = \psi_1 n$. This gives

$$N_a \cong \sqrt{\frac{B \sigma^2 n}{2 \psi_1 (n+2)}}. \quad (19)$$

which is also increasing in n , although at a lower rate. Thus, even under ‘‘CRTS’’ for the menu cost, the bundling of the goods pricing induces more frequent adjustments than in the case where the menu costs are dissociated, i.e. when $n = 1$.

We now study the invariant distribution of the sum of the squares of the price gaps $\|p\|^2 = \sum_{i=1}^n p_i(t)^2$ under the optimal policy. We will denote the density of the invariant distribution by $f(R)$ for $R \in [0, \bar{R}]$. This is interesting to study the response of a set of firms that are in steady state (i.e. in the invariant distribution) to an unexpected shock to their target which will displace the price gaps uniformly. In particular the study of how much mass is close to the boundary of inaction (so that after the ‘‘unexpected shock’’ they will decide to adjust) is one that has been identify as one of the key determinants to the impact effect of monetary policy shocks. We are interested in studying how this mass changes as we

⁶See Figures 1 and 2 in their paper. These authors group firms into 4 bins, according to the number of items sold (and recorded by the BLS), from 1 to 3 goods in the first bin to more than 7 goods in the fourth bin. They first measure the frequency of price changes at the good level, then compute the median frequency across the goods produced in the firm. Finally, they average these medians inside each of the 4 bins.

vary the number of goods n . There is an extensive literature on this topic, see for example Caballero and Engel (2007). The density of the invariant distribution has to solve the forward Kolmogorov equation and relevant boundary conditions.


PROPOSITION 5. The density $f(\cdot)$ of the invariant distribution of the sum of the squares of the price gaps R , for a given thresholds \bar{R} in the case of $n \geq 1$ products is for all $R \in [0, \bar{R}]$

$$\begin{aligned} f(R) &= \frac{1}{\bar{R}} [\log(\bar{R}) - \log(R)] \text{ if } n = 2, \text{ and} \\ f(R) &= (\bar{R})^{-\frac{n}{2}} \left(\frac{n}{n-2} \right) \left[(\bar{R})^{\frac{n}{2}-1} - (R)^{\frac{n}{2}-1} \right] \text{ otherwise.} \end{aligned} \quad (20)$$

So the density has a peak at $R = 0$, decreases in R , and reaches zero at \bar{R} . The shape depends on n . The density is convex in R for $n = 1, 2$ and $n = 3$, linear for $n = 4$, and concave for $n \geq 5$. This is intuitive, since the drift of the process for R increases linearly with n , hence the mass accumulates closer to the upper bound \bar{R} for higher n . Indeed as $n \rightarrow \infty$ the distribution converges to a uniform in $[0, \bar{R}]$. Proposition 5 makes clear also that the shape of the invariant density depends exclusively on n , the value of the other parameters, ψ, B, σ^2 only enters as determining \bar{R} , which only stretches the horizontal axis proportionally.

Consider the effect of decreasing the vector of price gaps by a constant $\delta > 0$ in all dimensions. The interpretation of this experiment, is the effect of an unexpected jump in the target price for all the goods. We want to find out the fraction of firms under the invariant that will adjust their prices. We will assume that from here on the price gap process remains the same, so the firms solved the problem stated above. For this we need the invariant distribution of p , not just R . We note that for each $R \leq \bar{R}$, the distribution of p is uniform on the n -dimensional sphere with radius \sqrt{R} . This is due to the symmetry of the distributions of the price gaps in each dimension, and its independence. Following the notation in Song and Gupta (1997) we use $U(n, 2)$ for the uniform distribution of p on the n -dimensional sphere, i.e. all the values of p with $\|p\| = 1$ have the same density. Now we obtain the fraction of prices p with $\|p\|^2 = R \leq \bar{R}$ that after the δ “shock” will be outside the set of inaction. They

Figure 1: $f(\cdot)$ density of invariant distribution of R , for various choices of n



are given by:

$$\|p\|^2 - 2\delta \left(\sum_{i=1}^n p_i \right) + n\delta^2 > \bar{R} \quad \text{or} \quad \frac{\sum_{i=1}^n p_i}{\sqrt{R}} \leq \frac{R - \bar{R}}{2\delta\sqrt{R}} + n\frac{\delta}{\sqrt{R}}. \quad (21)$$

Abusing notation let $u(x; n, 2)$ be the density of $U(n, 2)$ and define:

$$D(s) = \int_{\{x: \|x\|=1\}} \mathbf{I} \left\{ \sum_{i=1}^n p_i \leq s \right\} u(x_1, \dots, x_n; n, 2) dx_1 dx_2 \cdots dx_n.$$

Thus the fraction of firms that adjust after the unexpected increase in prices δ , denoted by is given by $A(\delta)$:

$$A(\delta) = \int_0^{\bar{R}} f(R) D \left(\frac{R - \bar{R}}{2\delta\sqrt{R}} + n\frac{\delta}{\sqrt{R}} \right) dR \approx -f'(\bar{R}) D \left(n\frac{\delta}{\sqrt{\bar{R}}} \right) \frac{\bar{R}^2}{2} = \frac{n}{2} D \left(n\frac{\delta}{\sqrt{\bar{R}}} \right), \quad (22)$$

where the last term is a an approximation of $A(\delta)$ using a first order expansion of the product

fD around \bar{R} , which will be accurate for small values of \bar{R} .

[TO FINISH THIS PART OF THE SECTION WE NEED TO DERIVE THE AVERAGE CHANGE ON PRICES OF THOSE THAT ADJUST]

PROPOSITION 6. Let t denote the time elapsed since the last price change. Let $J_\nu(\cdot)$ be the Bessel function of the first kind. The hazard rate for price changes is given by

$$h(t) = \sum_{r=1}^{\infty} \frac{\xi_{n,r}}{\sum_{s=1}^{\infty} \xi_{n,s} \exp\left(-\frac{q_{n,s}^2 \sigma^2}{2\bar{R}} t\right)} \frac{q_{n,r}^2 \sigma^2}{2\bar{R}} \exp\left(-\frac{q_{n,r}^2 \sigma^2}{2\bar{R}} t\right), \text{ where } \nu = \frac{n}{2} - 1,$$

$$\xi_{n,r} = \frac{1}{2^{\nu-1} \Gamma(\nu+1)} \frac{q_{n,r}^{\nu-1}}{J_{\nu+1}(q_{n,r})}, \text{ and } q_{nr} \text{ are the positive zeros of } J_\nu(\cdot),$$


which asymptotes to $\frac{\lim_{t \rightarrow \infty} h_n(t)}{\mathcal{T}(0)} = \frac{q_{n,1}^2}{2n} > \frac{(n-1)^2}{2n}$. (23)

In the proof we use results from probability theory on the first passage of time of a n brownian motion in a sphere center to the origin by [Ciesielski and Taylor \(1962\)](#) as well as characterization from the zeros of the Bessel function from [Qu and Wong \(1999\)](#) and [Hethcote \(1970\)](#).

Proposition 6 compares the asymptote of the hazard rate with the expected time until adjustment, which equals $\mathcal{T}(0) = \bar{R}/(n\sigma^2)$, as derived above. Notice that for a model with constant hazard rate these two quantities are the reciprocal of each other, i.e. the expected duration is the reciprocal of the hazard rate. We use this ratio, as a function of n as a measure of how close the model is to have constant hazard rates. We note that this ratio is exclusively a function of n . Indeed from the expression **Proposition 6**, it is immediate that the shape of the hazard rate function depends only on the number of products n . Changes in σ^2, B, ψ only stretch linearly the horizontal axis. More precisely, once keeping the expected time until adjustment $\mathcal{T}(0)$ fixed, the hazard rate is only a function of n .

Figure 2 plots the hazard rate function h for different choices of n keeping the expected time between price adjustment fixed at one. As **Proposition 6** shows the function h has an asymptote, which is increasing in the number of products n . Moreover, since the asymptote

Figure 2: Hazard rate of Price Adjustments for various choices of n



For each n the value of σ^2/\bar{R} is chosen so that the expected time elapsed between adjustments is one.

diverges to ∞ as n increases with no bound, the hazard rate converges to an inverted L shape, as the one for a model where adjustments are done exactly every $\mathcal{T}(0) = 1$ periods. To see this note that, defining $\tilde{R} \equiv R/\bar{R}$ and fixing the ratio $\sigma^2/\bar{R} = \mathcal{T}(0)/n$ so that for any n the expected time elapsed between price changes is $\mathcal{T}(0)$, we have:

$$d\tilde{R} = \mathcal{T}(0)dt + 2\sqrt{\tilde{R} \frac{\mathcal{T}(0)}{n}} dW \text{ for } \tilde{R} \in [0, 1]. \quad (24)$$

As $n \rightarrow \infty$ the process for the normalized size of the price gap \tilde{R} described in equation (24) converges to the deterministic one, in which case the hazard rate is zero between times 0 and below $\mathcal{T}(0)$ and ∞ precisely at $\mathcal{T}(0)$. For completeness, Table 1 computes the first zero for the relevant Bessel functions and the asymptotic hazard rate for several values of n .

[WE NEED TO CONCLUDE WITH THE IMPLICATIONS AS WE COMPARE WITH THE DATA]

Finally we discuss the distribution of price changes. This distribution can be described

Table 1: Limit hazard rates for various values of n

	number of products n									
	1	2	3	4	6	8	10	20	50	100
zeroes of $J_{\frac{n}{2}-1}(\cdot) : q_{n1}$	1.6.	2.4	3.1	3.8	5.1	6.4	7.6	13	30	56
$\frac{\text{Limit Hazard rate}}{\text{Expected duration}} : \frac{\lim_{t \rightarrow \infty} h_n(t, \bar{R})}{\mathcal{T}(0)}$	1.2	1.4	1.6	1.8	2.2	2.5	2.9	4.5	8.8	16

Note: for $n = 1$ and $n = 3$ the zeros are multiples of π , i.e. $q_{1,r} = (2(r-1) + 1)\pi/2$ and $q_{3,r} = r\pi$.

in terms of 2 parameters: the number of goods n , and the optimal threshold describing the size of the inaction set \bar{R} . The value of \bar{R} , as discussed above, depends on all the parameters. Since after an adjustment price gaps are set to zero, price changes coincide with the value of $p(\tau) \in \partial\mathcal{I} \subset \mathbb{R}^n$, the surface of an n -dimensional sphere of radius $\sqrt{\bar{R}}$. Let τ , be a time where p hits the boundary of the range of inaction, then given that each of the $p_i(t)$ when they are uncontrolled, are independently and identically normally distributed, price changes $\Delta p(\tau) = -p(\tau)$ are uniformly distributed in the n -dimensional surface of the sphere of radius $\sqrt{\bar{R}}$ ⁷. The next proposition characterizes the marginal distribution of price changes.

PROPOSITION 7. Let $\Delta p \in \partial\mathcal{I} \subset \mathbb{R}^n$ denote a price change for the n goods. The distribution of the price change of an individual good, i.e. the marginal distribution of $\Delta p_i \in [0, \sqrt{\bar{R}}]$ has density:

$$w(\Delta p_i) = \frac{1}{\text{Beta}\left(\frac{n-1}{2}, \frac{1}{2}\right) \sqrt{\bar{R}}} \left(1 - \left(\frac{\Delta p_i}{\sqrt{\bar{R}}}\right)^2\right)^{(n-3)/2} \quad (25)$$

where $\text{Beta}(\cdot, \cdot)$ denotes the Beta function. The standard deviation and kurtosis of the price changes, and expected value of the absolute value of price changes and its coefficient of

⁷The distribution of $\Delta p(\tau)$ is uniform in the surface of the sphere since the p.d.f. of a jointly normally distributed vector of n identical and independent normals, apart from a constant, is given by the exponential of a the square of an sphere with radius $\sqrt{\bar{R}}$.

variations are given by:

$$\begin{aligned} Std(\Delta p_i) &= \sqrt{\bar{R}/n}, \quad Kurt(\Delta p_i) = \frac{3n}{n+2}, \\ \mathbb{E}[|\Delta p_i|] &= \frac{\sqrt{\bar{R}}}{\frac{n-1}{2} Beta\left(\frac{n-1}{2}, \frac{1}{2}\right)}, \quad \frac{Std(|\Delta p_i|)}{\mathbb{E}(|\Delta p_i|)} = \sqrt{\left[\frac{n-1}{2} Beta\left(\frac{n-1}{2}, \frac{1}{2}\right)\right]^2 \frac{1}{n} - 1}. \end{aligned}$$

Moreover, as $n \rightarrow \infty$, the distribution of $\Delta p_i/Std(\Delta p_i)$ converges point-wise to a standard normal.

Using the previous proposition and the approximation for \bar{R} we obtain the following expression for the standard deviation of price changes:

$$Std(\Delta p_i) = \left(\frac{\sigma^2 \psi_1 2(n+2)}{B n^2}\right)^{1/4} \quad \text{and in the CRTS case } Std(\Delta p_i) = \left(\frac{\sigma^2 \psi_1 2(n+2)}{B n}\right)^{1/4},$$

where both expressions are decreasing in n . The expression for the kurtosis of the price changes shows that this statistic is an increasing function of n .

We can approximate some of the expressions in [Proposition 7](#) for statistics for $|\Delta p_i|$ involving the *Beta* function to obtain the following simpler expressions:⁸

$$\begin{aligned} \mathbb{E}[|\Delta p_i|] &\approx \sqrt{\bar{R}/n} \sqrt{2/\pi} \sqrt{1 + 1.1/(2n)} \quad \text{and} \\ \frac{Std(|\Delta p_i|)}{\mathbb{E}(|\Delta p_i|)} &\approx \sqrt{\frac{\pi}{2} \left(\frac{2n}{1.1 + 2n}\right) - 1}. \end{aligned}$$


The expression for the approximate value of $\mathbb{E}[|\Delta p_i|]$ is given by $Std(\Delta p_i)$ times a decreasing function of n . The expression for the approximate value of $Std(|\Delta p_i|)/\mathbb{E}(|\Delta p_i|)$ show that this statistic is an increasing function of n .

We note that the shape of the distribution h for price changes differs substantially for small values of n . For $n = 2$ is U-shaped, for $n = 3$ is uniform, for $n = 4$ it has the shape of

⁸ We note that error on the approximation error for $\mathbb{E}[|\Delta p_i|]$ and $Std(|\Delta p_i|)/\mathbb{E}(|\Delta p_i|)$ are smaller than 0.26% and 0.91%.

a half circle, and for $n \geq 6$ it has bell shape.⁹ Proposition 7 establishes that when $n \rightarrow \infty$ the distribution converges to a normal: this can be seen in Figure 3 by the comparison of the distribution for $n = 50$ and the p.d.f. of a normal distribution with standard deviation equal to $Std(\Delta p_i)$ for $n = 50$.

Figure 3: Density $w(\cdot)$ of the price changes for various choices of n



Parameter values: $B = 20$, $\sigma = 0.15$, $\psi_1 = 0.02$. Menu cost proportional to n . Solid lines are the p.d.f for h for different n . Circles denote the p.d.f. of a normal with standard deviation equal to that of Δp_i for $n = 50$.

Furthermore, from the expression in Proposition 7 the distribution of price changes x and of their absolute value $|x|$ depends on n and \bar{R} , any normalized statistics such as ratio of moments (kurtosis, skewness, etc) or a ratio of points in the c.d.f. depends exclusively on n . Indeed the kurtosis is given in Proposition 7, as $Kurtosis(\Delta p_i) = 3n/(2+n)$, and increasing concave function, starting at 1 and converging to 3. The next table computes several of these moments of interest, which has been computed for two scanner data sets by Midrigan (2009) to be compared with the prediction two menu cost models, one with one product and one with two.¹⁰

Comparing the statistics from Table 2 with the ones computed from scanner data by

⁹For n equal to 2, 3 and 4 one can grasp the shape of the distribution h from geometrical considerations,

Table 2: Statistics for price changes as function of number of products

statistics \ number of products n	1	2	3	4	5	6	10	20	50
$Std(\Delta p_i) / E(\Delta p_i)$	0	0.48	0.58	0.62	0.65	0.67	0.70	0.74	0.75
Kurtosis(Δp_i)	1.0	1.7	2.0	1.9	2.1	2.3	2.5	2.8	2.9
Fraction: $ \Delta p_i < \frac{1}{2}E(\Delta p_i)$	0	0.21	0.25	0.27	0.28	0.28	0.30	0.31	0.31
Fraction: $ \Delta p_i < \frac{1}{4}E(\Delta p_i)$	0	0.10	0.12	0.13	0.14	0.14	0.15	0.16	0.16

Δp_i denotes the log of the price change, and $|\Delta p_i|$ the absolute value of the log of price changes. They are computed the results in [Proposition 7](#). All statistics in the table depend exclusively on n . Kurtosis defined as the fourth moment relative to the square of the second.

[Midrigan \(2009\)](#) we conclude that a large value of n is required, on the order of 50. [THIS IS VERY SPECULATIVE, WE NEED TO EXPLAIN THE REASONS FOR THIS]

together with the fact the maximum of a density of a univariate normal is at one.

¹⁰The table in the [Midrigan \(2009\)](#) paper is Table 2b, Distribution of standardized price changes conditional on adjustments.

References

- Alvarez, Fernando E., Francesco Lippi, and Luigi Paciello. 2009. "Optimal price setting with information and menu costs." Mimeo - EIEF.
- Baccarin, Stefano. 2009. "Optimal impulse control for a multidimensional cash management system with generalized cost functions." *European Journal of Operational Research* 196 (1):198–206.
- Baudry, L., H. Le Bihan, P. Sevestre, and S. Tarrieu. 2007. "What do Thirteen Million Price Records have to Say about Consumer Price Rigidity?*" *Oxford Bulletin of Economics and Statistics* 69 (2):139–183.
- Bhattarai, Saroj and Raphael Schoenle. 2010. "Multiproduct Firms and Price-Setting: Theory and Evidence from U.S. Producer Prices." Working papers, Princeton University.
- Caballero, Ricardo J. and Eduardo M.R.A. Engel. 2007. "Price stickiness in Ss models: New interpretations of old results." *Journal of Monetary Economics* 54 (Supplemen):100–121.
- Cavallo, Alberto. 2009. "Scraped Data and Sticky Prices: Frequency, Hazards, and Synchronization ." Dissertation, Harvard University.
- Ciesielski, Z. and SJ Taylor. 1962. "First passage times and sojourn times for Brownian motion in space and the exact Hausdorff measure of the sample path." *Transactions of the American Mathematical Society* 103 (3):434–450.
- Dhyne, Emmanuel and Jerzy Konieczny. 2007. "Temporal Distribution of Price Changes: Staggering in the Large and Synchronization in the Small." Working Paper Series 01-07, Rimini Centre for Economic Analysis.
- Dixit, Avinash. 1991. "Analytical Approximations in Models of Hysteresis." *Review of Economic Studies* 58 (1):141–51.
- Dutta, Shantanu, Mark Bergen, Daniel Levy, and Robert Venable. 1999. "Menu Costs, posted prices and multiproduct retailers." *The Journal of Money, Credit and Banking* 31 (4):683–703.
- Golosov, Mikhail and Robert E. Jr. Lucas. 2007. "Menu Costs and Phillips Curves." *Journal of Political Economy* 115:171–199.
- Hethcote, H.W. 1970. "Bounds for zeros of some special functions." *Proceedings of the American Mathematical Society* :72–74.
- Karatzas, I. and S.E. Shreve. 1991. *Brownian motion and stochastic calculus*. Springer-Verlag.
- Karlin, S. and H.M. Taylor. 1999. *A second course in stochastic processes*. Academic press.
- Lach, S. and D. Tsiddon. 2007. "Small price changes and menu costs." *Managerial and Decision Economics* 28 (7):649–656.

- Lach, Saul and Daniel Tsiddon. 1992. “The Behavior of Prices and Inflation: An Empirical Analysis of Disaggregated Price Data.” *The Journal of Political Economy* 100 (2):349–389. URL <http://www.jstor.org/stable/2138611>.
- . 1996. “Staggering and Synchronization in Price-Setting: Evidence from Multiproduct Firms.” *The American Economic Review* 86 (5):1175–1196. URL <http://www.jstor.org/stable/2118285>.
- Midrigan, Virgiliu. 2007. “Menu Costs, Multi-Product Firms, and Aggregate Fluctuations.” CFS Working Paper Series 2007/13, Center for Financial Studies.
- . 2009. “Menu Costs, Multi-Product Firms, and Aggregate Fluctuations.” Working paper, NYU.
- Neiman, Brent. 2010. “Stickiness, synchronization, and passthrough in intrafirm trade prices.” *Journal of Monetary Economics* 57 (3):295–308.
- Qu, CK and R. Wong. 1999. “Best Possible Upper and Lower Bounds for the Zeros of the Bessel Function $J_\nu(x)$.” *Transactions of the American Mathematical Society* 351 (7):2833–2859.
- Song, D. and A. K. Gupta. 1997. “Lp-Norm Uniform Distribution.” *Proceedings of the American Mathematical Society* 125 (2):595–601.

A Proofs

Proof. (of [Proposition 2](#)) Notice that $v'(0) = \beta_1$ and that $v(0) = \beta_0$, so that we require $\beta_1 > 0$, which implies $\beta_0 > 0$. Moreover, if $\beta_1 > B/r$ then v is strictly increasing and strictly convex. If $\beta_1 = B/r$ then v is linear in R . If $0 < \beta_1 < B/r$, then v is strictly increasing at the origin, strictly concave, and it reaches its unique maximum at a finite value of R . Thus, a solution that satisfies smooth pasting requires that $0 < \beta_1 < B/r$, and the maximizer is \bar{R} . In this case, $R = 0$ achieves the minimum in the range $[0, \bar{R}]$. Finally, we require value matching at \bar{R} , i.e. $v(\bar{R}) = v(0) + \psi$. Let $\beta_i(\beta_1)$ be the solution of [equation \(15\)](#), as a function of β_1 . Note that for $0 < \beta_1 < B/r$, all the $\beta_i(\beta_1) < 0$ for $i \geq 2$ and are increasing in β_1 , converging to zero as β_1 goes to B/r . Smooth pasting can be written as

$$0 = v'(\bar{R}; \beta_1) \equiv \sum_{i=1}^{\infty} i \beta_i(\beta_1) \bar{R}^{i-1},$$

where we emphasize that all the β_i can be written as a function of β_1 . From the properties of the $\beta_i(\cdot)$ discussed above, it follows that we can write the unique solution of $0 = v'(\bar{\rho}(\beta_1); \beta_1)$ as an strictly increasing function of β_1 , i.e. $\bar{\rho}'(\beta_1) > 0$. Now we write value matching at \bar{R} which gives:

$$\psi = v(\bar{R}, \beta_1) - v(0, \beta_1) = v(\bar{R}, \beta_1) - \beta_0(\beta_1) = \sum_{i=1}^{\infty} \beta_i(\beta_1) \bar{R}^i.$$

We note that, given the properties of $\beta_i(\cdot)$ discussed above, for *any given* $R > 0$ we have: $v(R, \beta_1) - \beta_0(\beta_1)$ is strictly increasing in β_1 , as long as $0 < \beta_1 < B/r$. Thus, define

$$\Psi(\beta_1) = v(\bar{\rho}(\beta_1), \beta_1) - v(0, \beta_1) = \sum_{i=1}^{\infty} \beta_i(\beta_1) \bar{\rho}(\beta_1)^i .$$

From the properties discussed above we have that $\Psi(\beta_1)$ is strictly increasing in β_1 and that it ranges from 0 to ∞ as β_1 ranges from 0 to B/r . Thus Ψ is invertible. The solution of the problem is given by setting:

$$\beta_1(\psi) = \Psi^{-1}(\psi) \quad \text{and} \quad \bar{R}(\psi) = \bar{\rho}(\beta_1(\psi)) .$$

Now we show that the solution of the problem, as $r \downarrow 0$ satisfies:

$$\bar{R} = \sigma^2 q \left(\frac{\psi}{B \sigma^2}, n \right) .$$

for some increasing function $q(\cdot)$. As $r \downarrow 0$ solving the expected discounted problem is equivalent to solve the following steady state problem. The objective is to chose a threshold value of \bar{R} in order to minimize the sum of two terms. The first is expected number of adjustment times the cost per adjustment ψ from following the threshold policy. We denote this term by $N_a(\bar{R}; \sigma^2 n) \psi$. The second is the expected value of the deviations, times the cost of each deviation B . We denote this term by $e(\bar{R}; \sigma^2, n) B$. The minimization problem is then

$$\min_{\bar{R}} N_a(\bar{R}; \sigma^2 n) \psi + e(\bar{R}; \sigma^2, n) B$$

To be clear about the definition of this problem, denote $F(\cdot, n)$ the measure on the sample paths of n independent standard BM's, starting at zero at time zero. Let $\tau(\omega; \lambda, n)$ the first time that the sum of the squares of the BM's hits λ , and let $A(\lambda, n) = \{\omega : \sum_{i=1}^n W_i^2(\tau, \omega) = \lambda\}$ be the sample paths for which the sum of the squares hit λ . We have that

$$N_a(\bar{R}; \sigma^2 n) = \frac{1}{\int_{\omega \in A(\frac{\bar{R}}{\sigma^2}, n)} \tau(\omega; \frac{\bar{R}}{\sigma^2}, n) F(\omega, n) d\omega} = \frac{n \sigma^2}{\bar{R}}$$

where the last computation follow from [equation \(18\)](#) obtained in [Proposition 4](#).

$$e(\bar{R}; \sigma^2, n) \equiv \sigma^2 \int_{\omega \in A(\frac{\bar{R}}{\sigma^2}, n)} \frac{\int_0^{\tau(\omega; \frac{\bar{R}}{\sigma^2}, n)} \sum_{i=1}^n W_i^2(t, \omega) dt}{\tau(\omega; \frac{\bar{R}}{\sigma^2}, n)} F(\omega; n) d\omega$$

We note that since the n BM's have a normal finite distribution and are assumed to be independent, we have that

$$e(\bar{R}; \sigma^2, n) = \sigma^2 e \left(\frac{\bar{R}}{\sigma^2}; 1, n \right) .$$

We can write the minimization problem as:

$$B \min_{\bar{R}} \frac{n \sigma^2 \psi}{\bar{R} B} + \sigma^2 e \left(\frac{\bar{R}}{n \sigma^2}; 1, n \right) = \sigma^2 B \min_{\tilde{R}} \frac{n \psi}{\tilde{R} B \sigma^2} + e \left(\tilde{R}; 1, n \right)$$

where $\tilde{R} \equiv \bar{R}/\sigma^2$. Let the solution of the transformed problem be $\tilde{R} = q \left(\frac{\psi}{B \sigma^2}, n \right)$ then the solution of the original problem is:

$$\bar{R} = \sigma^2 q \left(\frac{\psi}{B \sigma^2}, n \right) .$$

That the function $q(\cdot)$ is strictly increasing follows from the previous result and since ψ is an argument of this function. Q.E.D.

Proof. (of [Proposition 3](#)) We show that V so constructed has the following properties:

1. it only depends on the absolute value of the prices, since for all $p \in \mathbb{R}^n$:

$$v \left(\sum_{i=1}^n p_i^2 \right) = v \left(\sum_{i=1}^n |p_i|^2 \right) .$$

for all $p \in \mathbb{R}^n$,

2. The range of inaction is given by $\mathcal{I} = \{p \in \mathbb{R}^n \mid \sum_{i=1}^n p_i^2 \leq \bar{R}\}$.
3. It solves the ODE given by [equation \(3\)](#). This can be seen by computing:

$$V_i(p) = v' \left(\sum_{i=1}^n p_i^2 \right) 2p_i \quad \text{and} \quad V_{ii}(p) = v'' \left(\sum_{i=1}^n p_i^2 \right) (2p_i)^2 + v' \left(\sum_{i=1}^n p_i^2 \right) 2 ,$$

replacing this into the ODE [equation \(3\)](#) we obtain the ODE [equation \(10\)](#), which v solves by hypothesis.

4. It satisfies value matching [equation \(4\)](#), which is immediate since it satisfied the value matching condition for v given in [equation \(11\)](#).
5. it satisfies smooth pasting [equation \(6\)](#). Using the form of the solution for v , namely:

$$V_i(p) = v' \left(\sum_{i=1}^n p_i^2 \right) 2 p_i = \sum_{j=1}^n \beta_j j \left(\sum_{k=1}^n p_k^2 \right)^{j-1} 2 p_i$$

Using that v satisfies smooth pasting we have:

$$0 = \sum_{j=1}^n \beta_j j \left(\sum_{k=1}^n p_k^2 \right)^{j-1}$$

for any p with $\sum_{k=1}^n p_k^2 = \bar{R}$, which establishes that $V_i(p) = 0$ for all $i = 1, \dots, n$ and for any $p \in \partial \mathcal{I}$.

6. It satisfies optimality of the origin as return point, as given by [equation \(5\)](#). Direct computation gives:

$$V_i(p) = v' \left(\sum_{i=1}^n p_i^2 \right) 2 p_i = \sum_{j=1}^n \beta_j j \left(\sum_{k=1}^n p_k^2 \right)^{j-1} 2 p_i$$

which equals zero when evaluated at $p = 0$. Notice also that

$$V_{ii}(0) = 2 \beta_1 > 0 \text{ for all } i = 1, \dots, n \text{ and } V_{ij}(0) = 0 ,$$

thus, $p = 0$ is a local minimum.

Finally, a function V with these properties is a strong solution to the variational inequality of the problem, and hence it is the value function. QED

Proof. (of [Proposition 5](#)) The forward Kolmogorov equation is:

$$0 = \frac{1}{2} \frac{\partial^2}{\partial R^2} \left(\left[2\sigma\sqrt{R} \right]^2 f(R) \right) - \frac{\partial}{\partial R} (n\sigma^2 f(R)) \text{ for } R \in (0, \bar{R}) , \quad (26)$$

with boundary conditions:

$$1 = \int_0^{\bar{R}} f(R) dR \text{ and } f(\bar{R}) = 0 . \quad (27)$$

The first boundary conditions ensures that f is a density. The second is implied by the fact that when the process reaches \bar{R} it is return to the origin, so the mass escape from these points. [Equation \(27\)](#) implies the second order ODE: $f'(R)(\frac{n}{2} - 2) = Rf''(R)$. The solution of this ODE for $n \neq 2$ is $f(R) = A_1 R^{n/2-1} + A_0$ for two constants A_0, A_1 to be determine using the boundary conditions [equation \(27\)](#):

$$\begin{aligned} 0 &= A_1(\bar{R})^{n/2-1} + A_0 , \\ 1 &= \frac{A_1}{n/2}(\bar{R})^{n/2} + A_0\bar{R} . \end{aligned}$$

For $n = 2$ the solution is $f(R) = -A_1 \log(R) + A_0$ subject to the analogous conditions. Solving for the coefficients A_0, A_1 gives the desired expressions.

Proof. (of [Proposition 6](#)) Let τ be the stopping time defined by the first time where the sum of the square of the price gaps vector $\|p(\tau)\|^2$ reaches the critical value \bar{R} , starting at the origin at time zero, i.e. starting at $\|p(0)\| = 0$. Let $S_n(t, \bar{R})$ be the probability distribution for stopping times $\tau \geq t$, alternatively let $S_n(\cdot, \bar{R})$ be the survival function. [Theorem 2 Ciesielski and Taylor \(1962\)](#) shows that for $n \geq 1$:

$$S_n(t, \bar{R}) = \sum_{r=1}^{\infty} \xi_{n,r} \exp \left(-\frac{q_{n,r}^2}{2\bar{R}} \sigma^2 t \right) , \text{ where } \xi_{n,r} = \frac{1}{2^{\nu-1}\Gamma(\nu+1)} \frac{q_{n,r}^{\nu-1}}{J_{\nu+1}(q_{n,r})} . \quad (28)$$

where $J_\nu(z)$ is the Bessel function of the first kind, where $\nu = (n-2)/2$, where q_{nr} are the positive zeros of the Bessel function $J_\nu(z)$, index in ascending order according to r , and where

Γ is the gamma function. The hazard rate is then given by:

$$h_n(t, \bar{R}) = -\frac{1}{S_n(t, \bar{R})} \frac{\partial S_n(t, \bar{R})}{\partial t}, \text{ with asymptote } \lim_{t \rightarrow \infty} h_n(t, \bar{R}) = \frac{q_{n,1}^2 \sigma^2}{2 \bar{R}}. \quad (29)$$

As shown by [Qu and Wong \(1999\)](#), the zeroes of the Bessel function $q_{n,r}$ satisfy for $n > 2$ the following inequalities:

$$\left(\frac{n}{2} - 1\right) - \frac{a_r}{2^{1/3}} \left(\frac{n}{2} - 1\right)^{1/3} < q_{n,r} < \left(\frac{n}{2} - 1\right) - \frac{a_r}{2^{1/3}} \left(\frac{n}{2} - 1\right)^{1/3} + \frac{3}{10} a_r^2 \frac{2^{1/3}}{\left(\frac{n}{2} - 1\right)^{1/3}} \quad (30)$$

where a_r are the first negative zero of the Airy function. For instance $a_1 \approx -2.33811$, giving a tight bound for the first zero $q_{n,1}$, which determines the asymptote of the hazard rate. A related simpler lower bound given by [Hethcote \(1970\)](#) for $n \geq 2$ is

$$q_{n,r}^2 > \left(r - \frac{1}{4}\right)^2 \pi^2 + \left(\frac{n}{2} - 1\right)^2. \quad (31)$$

Proof. (of [Proposition 7](#)) We first establish the following Lemma. Lemma. Let z be distributed uniformly on the surface in the surface of the n -dimensional sphere of radius one. We use x for the projection of z in any of the dimension, so $z_i = x \in [-1, 1]$. The marginal distribution of $x = z_i$ has density:

$$\begin{aligned} f_n(x) &= \int_0^\infty \frac{s^{(n-3)/2} e^{-s/2}}{2^{(n-1)/2} \Gamma[(n-1)/2]} \frac{e^{-sx^2/[2(1-x^2)]}}{\sqrt{2\pi}} \frac{s^{1/2}}{(1-x^2)^{3/2}} ds \\ &= \frac{\Gamma(n/2)}{\Gamma(1/2) \Gamma[(n-1)/2]} (1-x^2)^{(n-3)/2} \end{aligned} \quad (32)$$

where the Γ function makes the density integrate to one. This lemma is an application of Theorem 2.1, part 1 in [Song and Gupta \(1997\)](#), setting $p = 2$, so it is euclidian norm, and $k = 1$ so it is the marginal of one dimension. We give a simpler proof below.

Now we consider the case where the sphere has radius different from one. Let $p \in \partial\mathcal{I}$, then

$$p = \frac{p}{\sum_{i=1}^n p_i^2} \bar{R} = \frac{p}{\sqrt{\sum_{i=1}^n p_i^2}} \sqrt{\bar{R}} = z \sqrt{\bar{R}}$$

where z is uniformly distributed in the n dimensional sphere of radius one. Thus each p_i has the same distribution than $x \sqrt{\bar{R}}$. Using the change of variable formula we obtained the required result.

Part 2 of Theorem 2.1 in [Song and Gupta \(1997\)](#) shows that if x the marginal of a uniform distributed vector in the surface of the n -dimensional sphere, then x^2 is distributed as a $Beta(\frac{1}{2}, n-12)$. If y is distributed as a $Beta(\alpha, \beta)$ then it has $\mathbb{E}(y) = \alpha/(\alpha + \beta)$ and $\mathbb{E}(y^2) = (\alpha+1)/(\alpha + \beta + 1)\mathbb{E}(y)$. Using these expressions for $\alpha = 1/2$ and $\beta = n/2$ we obtain the results for the standard deviation of Δp_i and its kurtosis. For the expected value of the

absolute value of price changes we note that

$$\begin{aligned}
\mathbb{E}[|\Delta p_i|] &= 2 \int_0^{\sqrt{R}} \Delta p_i w(\Delta p_i) d\Delta p_i \\
&= \frac{2}{\text{Beta}\left(\frac{n-1}{2}, \frac{1}{2}\right)} \int_0^{\sqrt{R}} \Delta p_i \left(1 - \left(\frac{\Delta p_i}{\sqrt{R}}\right)^2\right)^{(n-3)/2} d\Delta p_i \\
&= \frac{\sqrt{R}}{\frac{n-1}{2} \text{Beta}\left(\frac{n-1}{2}, \frac{1}{2}\right)}
\end{aligned}$$

where the second line uses the form of h and the last line uses that the following result:

$$\int_a^b x (1-x^2)^{(n-3)/2} dx = \frac{(1-x^2)^{(n-1)/2}}{1-n} \Big|_a^b.$$

Then we have, using the fundamental property of the Gamma function

$$\frac{1}{\frac{n-1}{2} \text{Beta}\left(\frac{n-1}{2}, \frac{1}{2}\right)} = \frac{\Gamma\left(\frac{n}{2}\right)}{\frac{n-1}{2} \Gamma\left(\frac{n-1}{2}\right) \Gamma(1/2)} = \frac{\Gamma\left(\frac{n}{2}\right)}{\Gamma\left(\frac{n+1}{2}\right) \Gamma(1/2)}$$

Thus

$$\mathbb{E}[|\Delta p_i|] = \frac{\sqrt{R}}{\frac{n-1}{2} \text{Beta}\left(\frac{n-1}{2}, \frac{1}{2}\right)} = \sqrt{R} \frac{\Gamma\left(\frac{n}{2}\right)}{\Gamma\left(\frac{n+1}{2}\right) \Gamma(1/2)}$$

We can approximate these ratio of Gamma functions as

$$\frac{\Gamma\left(\frac{n}{2}\right)}{\Gamma\left(\frac{n+1}{2}\right) \Gamma(1/2)} \approx \sqrt{\frac{2}{\pi}} \frac{\sqrt{n+1/2}}{n}$$

from where we obtain our expression.

For $Std(|\Delta p_i|) / \mathbb{E}(|\Delta p_i|)$ we use that, given the symmetry around zero we have:

$$\begin{aligned}
Std(|\Delta p_i|) / \mathbb{E}(|\Delta p_i|) &= \sqrt{\frac{\mathbb{E}[\Delta p_i^2]}{\mathbb{E}[|\Delta p_i|]^2} - 1} = \sqrt{\left(\frac{Std(\Delta p_i)}{\mathbb{E}[|\Delta p_i|]}\right)^2 - 1} \\
&= \sqrt{\left(\frac{\frac{n-1}{2} \text{Beta}\left(\frac{n-1}{2}, \frac{1}{2}\right)}{\sqrt{n}}\right)^2 - 1} \approx \sqrt{\frac{\pi}{2} \left(\frac{2n}{1+2n}\right) - 1}
\end{aligned}$$

For the convergence of $\Delta p_i / Std(\Delta p_i)$ to a normal, we show that $y = x^2 n$ converges to a chi-square distribution with 1 d.o.f., where x is the marginal of a uniform distribution in the surface of the n -dimensional sphere. The p.d.f of $y \in [0, n]$, the square of the standardized x , is

$$\frac{\Gamma\left(\frac{n}{2}\right)}{n \Gamma\left(\frac{n-1}{2}\right) \Gamma\left(\frac{1}{2}\right)} \left(1 - \left(\frac{y}{n}\right)\right)^{(n-3)/2} \left(\frac{y}{n}\right)^{-1/2},$$

and the p.d.f. of a chi-square with 1 d.o.f. is

$$\frac{\exp(-y/2) y^{-1/2}}{\sqrt{2} \Gamma(\frac{1}{2})}.$$

Then, fixing y , taking logs in the ratio of the two p.d.f.'s, and taking the limit as $n \rightarrow \infty$, using that

$$\frac{\Gamma(\frac{n}{2}) \sqrt{2}}{\Gamma(\frac{n-1}{2}) \sqrt{n}} \rightarrow 1 \text{ as } n \rightarrow \infty.$$

we obtain that the ratio of the two p.d.f.'s converges to one.

Q.E.D.

Derivation of the approximation for \bar{R} . The smooth pasting and value matching conditions gives a system of two equations in two unknowns,

$$\begin{aligned} 0 &= v'(\bar{R}) = \beta_1 + 2\beta_2\bar{R} \\ \psi &= v(\bar{R}) - v(0) = \beta_1\bar{R} + \beta_2\bar{R}^2, \end{aligned}$$

where $\beta_2 = (r\beta_1 - B)/(2\sigma^2(n+2))$. We replace \bar{R} and β_2 to obtain the following quadratic solution for β_1 :

$$-\beta_1^2 - r\beta_1 \left(\frac{2\psi}{\sigma^2(n+2)} \right) + \frac{B\psi}{\sigma^2(n+2)} = 0$$

taking the positive root and replacing it back to solve for \bar{R} yields:

$$\bar{R} = \frac{2\psi}{-\frac{r\psi}{\sigma^2(n+2)} + \sqrt{\left(\frac{r\psi}{\sigma^2(n+2)}\right)^2 + \frac{2B\psi}{\sigma^2(n+2)}}$$

Taking r to zero in this expression we obtain [equation \(16\)](#).


B Numerical accuracy of the approximation

In this section we present some evidence on the numerical accuracy of the approximation. We compare the value of \bar{R} obtained from the quadratic approximation to v described above, with what we call the ‘‘exact’’ solution, which is the numerical solution using up to 30 terms for β_i in its the expansion.

The approximation are closer for smaller values of σ and ψ , which we regard as more realistic.


The next figure shows the value of $N_a(n)$ for various n when the menu cost are constant returns to scale, so $\psi = \psi_1 n$ using the approximation and using the ‘‘exact’’ expression.

Figure 4: Ratio of \bar{R} 's and of $v(0)$'s for the approximation relative to the “exact” solution



Note: parameter values are $B = 20$, $\sigma = 0.25$, $\psi_1 = 0.03$ and $r = 0.03$.

Figure 5: Frequency of adjustment N_a for the CRTS $\psi_1 n$ and constant ψ .



Note: parameter values are $B = 20$, $\sigma = 0.20$, $\psi_1 = 0.03$ and $r = 0.03$.

C Approximating the Profit Function

Consider the expression for the rate of profits of a multiproduct firm. The marginal cost for each of the products is C_i . The demand system is given by the sum of n independent demands, with own price elasticity given by η . The parameter A_i is the intercept, in logs, of the demand for the i -th product. Given the constant elasticity of demand and constant marginal cost, the frictionless optimal price for the monopolist is a multiple of the marginal cost, and independent of A_i . To keep the n goods symmetric we will assume that C_i and A_i are perfectly correlated, so that when cost are high, and hence frictionless prices are high, demand is also high. In this way we can keep the share of profits coming to each of the n goods comparable, even if cost differ significantly.

We write the total profits per product

$$\Pi(P_1, \dots, P_n, C_1, \dots, C_n, A_1, \dots, A_n) \equiv \sum_{i=1}^n \Pi(P_i, C_i, A_i) = \sum_{i=1}^n A_i P_i^{-\eta} (P_i - C_i)$$

Let $P_i^* = \arg \max_P \Pi(P, C_i, A_i)$. Assuming that

$$A_i = A (C_i)^{\eta-1},$$

we obtain that profits, relative to the maximized profits, can be written as

$$\begin{aligned} & \frac{\Pi(P_1, \dots, P_n, C_1, \dots, C_n, A_1, \dots, A_n) - \Pi(P_1^*, \dots, P_n^*, C_1, \dots, C_n, A_1, \dots, A_n)}{\Pi(P_1^*, \dots, P_n^*, C_1, \dots, C_n, A_1, \dots, A_n)} \\ &= B \sum_{i=1}^n \left(\frac{P_i - P_i^*}{P_i^*} \right)^2 + o \left(\sum_{i=1}^n \left(\frac{P_i - P_i^*}{P_i^*} \right)^2 \right) \end{aligned}$$

where $B = \frac{(\eta-1)\eta}{2n}$.

To obtain the quadratic expression above we write a second order expansion of the profits, divide both sides by the maximized total profits, and complete elasticities:

$$\begin{aligned} & \frac{\Pi(P_1, \dots, P_n, C_1, \dots, C_n, A_1, \dots, A_n)}{\Pi(P_1^*, \dots, P_n^*, C_1, \dots, C_n, A_1, \dots, A_n)} \\ &= 1 + \sum_{i=1}^n \frac{1}{\Pi(P_1^*, \dots, P_n^*, C_1, \dots, C_n, A_1, \dots, A_n)} \frac{\partial}{\partial P_i} \Pi(P_i, C_i, A_i) \Big|_{P_i^*} P_i^* \left(\frac{P_i - P_i^*}{P_i^*} \right) \\ &+ \frac{1}{2} \sum_{i=1}^n \frac{1}{\Pi(P_1^*, \dots, P_n^*, C_1, \dots, C_n, A_1, \dots, A_n)} \frac{\partial^2}{\partial P_i^2} \Pi(P_i, C_i) \Big|_{P_i^*} (P_i^*)^2 \left(\frac{P_i - P_i^*}{P_i^*} \right)^2 \end{aligned}$$

Computing the derivatives for our functional forms:

$$\begin{aligned}\frac{\partial}{\partial P_i} \Pi(P_i, C_i, A_i) &= A_i P^{-\eta} \left(-\eta \left(\frac{P_i - C_i}{P_i} \right) + 1 \right) \\ \frac{\partial^2}{\partial P_i^2} \Pi(P_i, C_i) &= -A_i P^{-\eta} \eta \frac{1}{P_i} \left(-\eta \left(\frac{P_i - C_i}{P_i} \right) + 1 \right) - A_i P^{-\eta} \eta \left(\frac{C_i}{P_i^2} \right)\end{aligned}$$

We have the standard result of a constant mark-up:

$$P^* = \frac{\eta}{\eta - 1} C_i \implies \left(-\eta \left(\frac{P_i - C_i}{P_i} \right) + 1 \right) = \left(-\eta \left(\frac{\eta C_i - C_i(\eta - 1)}{\eta C_i} \right) + 1 \right) = 0,$$

and the maximized value of profits given by

$$\Pi(P_i^*, C_i, A_i) = A_i C_i^{-\eta} \left(\frac{\eta}{\eta - 1} \right)^{-\eta} C_i \left(\frac{1}{\eta - 1} \right) = A_i C_i^{1-\eta} \left(\frac{\eta}{\eta - 1} \right)^{-\eta} \left(\frac{1}{\eta - 1} \right).$$

Hence the first and second derivatives, evaluated at the optimal prices are:

$$\begin{aligned}\frac{\partial}{\partial P_i} \Pi(P_i, C_i, A_i) |_{P^*} &= 0 \\ \frac{\partial^2}{\partial P_i^2} \Pi(P_i, C_i, A_i) |_{P^*} &= -A_i P^{*-\eta} \eta \frac{C_i}{P_i^{*2}} = -A_i \left(C_i \frac{\eta}{\eta - 1} \right)^{-\eta} \frac{\eta C_i}{P_i^{*2}}\end{aligned}$$

and

$$\frac{1}{\Pi(P_i^*, C_i, A_i)} \frac{\partial^2}{\partial P_i^2} \Pi(P_i, C_i, A_i) |_{P^*} (P_i^*)^2 = -\frac{A_i \left(C_i \frac{\eta}{\eta - 1} \right)^{-\eta} \eta C_i}{A_i C_i^{1-\eta} \left(\frac{\eta}{\eta - 1} \right)^{-\eta} \left(\frac{1}{\eta - 1} \right)} = -(\eta - 1) \eta$$

Thus the expansion can be written as:

$$\begin{aligned}& \frac{\Pi(P_1, \dots, P_n, C_1, \dots, C_n, A_1, \dots, A_n)}{\Pi(P_1^*, \dots, P_n^*, C_1, \dots, C_n, A_1, \dots, A_n)} \\ &= 1 + \frac{1}{2} \sum_{i=1}^n \frac{1}{\Pi(P_1^*, \dots, P_n^*, C_1, \dots, C_n, A_1, \dots, A_n)} \frac{\partial^2}{\partial P_i^2} \Pi(P_i, C_i) |_{P_i^*} (P_i^*)^2 \left(\frac{P_i - P_i^*}{P_i^*} \right)^2 \\ &= 1 - \frac{1}{2} \sum_{i=1}^n \frac{\Pi(P_i^*, C_i, A_i)}{\Pi(P_1^*, \dots, P_n^*, C_1, \dots, C_n, A_1, \dots, A_n)} (\eta - 1) \eta \left(\frac{P_i - P_i^*}{P_i^*} \right)^2\end{aligned}$$

Using the assumption that $A_i = A (C_i)^{\eta-1}$ we have that

$$\frac{\Pi(P_i^*, C_i, A_i)}{\Pi(P_1^*, \dots, P_n^*, C_1, \dots, C_n, A_1, \dots, A_n)} = \frac{A_i C_i^{1-\eta}}{\sum_{j=1}^n A_j C_j^{1-\eta}} = \frac{1}{n},$$

and hence the expression for B is:

$$B = \frac{(\eta - 1)\eta}{2n}.$$