

CLUSTER SAMPLES AND CLUSTERING

Jeff Wooldridge
Michigan State University
LABOUR Lectures, EIEF
October 18-19, 2011

1. The Linear Model with Cluster Effects
2. Cluster-Robust Inference with Large Group Sizes
3. Cluster Samples with Unit-Specific Panel Data
4. Clustering and Stratification
5. Two-Way Clustering

1. The Linear Model with Cluster Effects

- For each group or cluster g , let $\{(y_{gm}, \mathbf{x}_g, \mathbf{z}_{gm}) : m = 1, \dots, M_g\}$ be the observable data, where M_g is the number of units in cluster or group g , y_{gm} is a scalar response, \mathbf{x}_g is a $1 \times K$ vector containing explanatory variables that vary only at the cluster or group level, and \mathbf{z}_{gm} is a $1 \times L$ vector of covariates that vary within (as well as across) groups.

- Without a cluster identifier, a cluster sample looks like a cross section data set. Statistically, the key difference is that the sample of clusters has been drawn from a “large” population of clusters.
- The clusters are assumed to be independent of each other, but outcomes within a cluster should be allowed to be correlated.

- An example is randomly drawing fourth-grade classrooms from a large population of classrooms (say, in the state of Michigan). Each class is a cluster and the students within a class are the individual units. Or we draw industries and then we have firms within an industry. Or we draw hospitals and then we have patients within a hospital.
- If higher-level explanatory variables are included in a model, we should consider the data as a cluster sample at the higher level to ensure valid inference.

- The linear model with an additive error is

$$y_{gm} = \alpha + \mathbf{x}_g\boldsymbol{\beta} + \mathbf{z}_{gm}\boldsymbol{\gamma} + v_{gm} \quad (1.1)$$

for $m = 1, \dots, M_g, g = 1, \dots, G$.

- The observations are independent across g (group or cluster).

- Key questions:

(1) Are we primarily interested in β (group-level coefficients) or γ (individual-level coefficients)?

(2) Does v_{gm} contain a common group effect, as in

$$v_{gm} = c_g + u_{gm}, m = 1, \dots, M_g, \quad (1.2)$$

where c_g is an unobserved group (cluster) effect and u_{gm} is the idiosyncratic component? (Act as if it does.)

(3) Are the regressors $(\mathbf{x}_g, \mathbf{z}_{gm})$ appropriately exogenous?

(4) How big are the group sizes (M_g) and number of groups (G)? For now, we are assuming “large” G and “small” M_g , but we cannot give specific values.

- The theory with $G \rightarrow \infty$ and fixed group sizes, M_g , is well developed [White (1984), Arellano (1987)].
- How should one use the theory? If we assume

$$E(v_{gm} | \mathbf{x}_g, \mathbf{z}_{gm}) = 0 \tag{1.3}$$

then pooled OLS estimator of y_{gm} on

$1, \mathbf{x}_g, \mathbf{z}_{gm}, m = 1, \dots, M_g; g = 1, \dots, G$, is consistent for $\boldsymbol{\lambda} \equiv (\alpha, \boldsymbol{\beta}', \boldsymbol{\gamma}')'$

(as $G \rightarrow \infty$ with M_g fixed) and \sqrt{G} -asymptotically normal.

- Robust variance matrix is needed to account for correlation within clusters or heteroskedasticity in $Var(v_{gm}|\mathbf{x}_g, \mathbf{z}_{gm})$, or both. Write \mathbf{W}_g as the $M_g \times (1 + K + L)$ matrix of all regressors for group g . Then the $(1 + K + L) \times (1 + K + L)$ variance matrix estimator is

$$\left(\sum_{g=1}^G \mathbf{W}'_g \mathbf{W}_g \right)^{-1} \left(\sum_{g=1}^G \mathbf{W}'_g \hat{\mathbf{v}}_g \hat{\mathbf{v}}'_g \mathbf{W}_g \right) \left(\sum_{g=1}^G \mathbf{W}'_g \mathbf{W}_g \right)^{-1}, \quad (1.4)$$

where $\hat{\mathbf{v}}_g$ is the $M_g \times 1$ vector of pooled OLS residuals for group g .

This “sandwich” estimator is now computed routinely using “cluster” options.

- Sometimes an adjustment is made, such as multiplying by $G/(G - 1)$.

- In Stata, used “cluster” option with standard regression command:

```
reg y x1 ... xK z1 ... zL, cluster(clusterid)
```

- These standard errors are, as in the panel data case, robust to unknown heteroskedasticity, too.
- Structure of asymptotic variance is identical to panel data case (because $G \rightarrow \infty$ plays the role of $N \rightarrow \infty$ and M_g fixed is like fixed T in panel data case).
- Cluster samples are usually “unbalanced,” that is, the M_g vary across g .

- Generalized Least Squares: Strengthen the exogeneity assumption to

$$E(v_{gm}|\mathbf{x}_g, \mathbf{Z}_g) = 0, m = 1, \dots, M_g; g = 1, \dots, G, \quad (1.5)$$

where \mathbf{Z}_g is the $M_g \times L$ matrix of unit-specific covariates. Condition (1.5) is “strict exogeneity” for cluster samples (without a time dimension).

- If \mathbf{z}_{gm} includes only unit-specific variables, (1.5) rules out “peer effects.” But one can include measures of peers in \mathbf{z}_{gm} – for example, the fraction of friends living in poverty or living with only one parent.

- Full RE approach: the $M_g \times M_g$ variance-covariance matrix of $\mathbf{v}_g = (v_{g1}, v_{g2}, \dots, v_{g,M_g})'$ has the “random effects” form,

$$\text{Var}(\mathbf{v}_g) = \sigma_c^2 \mathbf{j}'_{M_g} \mathbf{j}_{M_g} + \sigma_u^2 \mathbf{I}_{M_g}, \quad (1.6)$$

where \mathbf{j}_{M_g} is the $M_g \times 1$ vector of ones and \mathbf{I}_{M_g} is the $M_g \times M_g$ identity matrix.

- The usual assumptions include the “system homoskedasticity” assumption,

$$\text{Var}(\mathbf{v}_g | \mathbf{x}_g, \mathbf{Z}_g) = \text{Var}(\mathbf{v}_g). \quad (1.7)$$

- The random effects estimator $\hat{\lambda}_{RE}$ is asymptotically more efficient than pooled OLS under (1.5), (1.6), and (1.7) as $G \rightarrow \infty$ with the M_g fixed. The RE estimates and test statistics for cluster samples are computed routinely by popular software packages (sometimes by making it look like a panel data set).

- An important point is often overlooked: one can, and in many cases should, make RE inference completely robust to an unknown form of $Var(\mathbf{v}_g|\mathbf{x}_g, \mathbf{Z}_g)$ even in the cluster sampling case.
- The motivation for using the usual RE estimator when $Var(\mathbf{v}_g|\mathbf{x}_g, \mathbf{Z}_g)$ does not have the RE structure is the same as that for GEE: the RE estimator may be more efficient than POLS.

- Example: Random coefficient model,

$$y_{gm} = \alpha + \mathbf{x}_g \boldsymbol{\beta} + \mathbf{z}_{gm} \boldsymbol{\gamma}_g + v_{gm}. \quad (1.8)$$

By estimating a standard random effects model that assumes common slopes $\boldsymbol{\gamma}$, we effectively include $\mathbf{z}_{gm}(\boldsymbol{\gamma}_g - \boldsymbol{\gamma})$ in the idiosyncratic error:

$$y_{gm} = \alpha + \mathbf{x}_g \boldsymbol{\beta} + \mathbf{z}_{gm} \boldsymbol{\gamma} + c_g + [u_{gm} + \mathbf{z}_{gm}(\boldsymbol{\gamma}_g - \boldsymbol{\gamma})]$$

- The usual RE transformation does not remove the correlation across errors due to $\mathbf{z}_{gm}(\boldsymbol{\gamma}_g - \boldsymbol{\gamma})$, and the conditional correlation depends on \mathbf{Z}_g in general.

- If only γ is of interest, fixed effects is attractive. Namely, apply pooled OLS to the equation with group means removed:

$$y_{gm} - \bar{y}_g = (\mathbf{z}_{gm} - \bar{\mathbf{z}}_g)\boldsymbol{\gamma} + u_{gm} - \bar{u}_g. \quad (1.9)$$

- FE allows arbitrary correlation between c_g and $\{\mathbf{z}_{gm} : m = 1, \dots, M_g\}$.

• Can be important to allow $Var(\mathbf{u}_g|\mathbf{Z}_g)$ to have arbitrary form, including within-group correlation and heteroskedasticity. Using the argument for the panel data case, FE can consistently estimate the average effect in the random coefficient case. But $(\mathbf{z}_{gm} - \bar{\mathbf{z}}_g)(\gamma_g - \gamma)$ appears in the error term:

$$y_{gm} - \bar{y}_g = (\mathbf{z}_{gm} - \bar{\mathbf{z}}_g)\boldsymbol{\gamma} + (u_{gm} - \bar{u}_g) + (\mathbf{z}_{gm} - \bar{\mathbf{z}}_g)(\gamma_g - \gamma)$$

- A fully robust variance matrix estimator of $\hat{\boldsymbol{\gamma}}_{FE}$ is

$$\left(\sum_{g=1}^G \ddot{\mathbf{Z}}_g' \ddot{\mathbf{Z}}_g \right)^{-1} \left(\sum_{g=1}^G \ddot{\mathbf{Z}}_g' \hat{\mathbf{u}}_g \hat{\mathbf{u}}_g' \ddot{\mathbf{Z}}_g \right) \left(\sum_{g=1}^G \ddot{\mathbf{Z}}_g' \ddot{\mathbf{Z}}_g \right)^{-1}, \quad (1.10)$$

where $\ddot{\mathbf{Z}}_g$ is the matrix of within-group deviations from means and $\hat{\mathbf{u}}_g$ is the $M_g \times 1$ vector of fixed effects residuals. This estimator is justified with large- G asymptotics.

- Can also use pooled OLS or RE on

$$y_{gm} = \alpha + \mathbf{x}_g\boldsymbol{\beta} + \mathbf{z}_{gm}\boldsymbol{\gamma} + \bar{\mathbf{z}}_g\boldsymbol{\xi} + e_{gm}, \quad (1.11)$$

which allows inclusion of \mathbf{x}_g and a simple test of $H_0 : \boldsymbol{\xi} = \mathbf{0}$. Again, fully robust inference.

- POLS and RE of (1.11) both give the FE estimate of $\boldsymbol{\gamma}$.
- Example: Estimating the Salary-Benefits Tradeoff for Elementary School Teachers in Michigan.
- Clusters are school districts. Units are schools within a district.

. des

Contains data from C:\mitbook1_2e\statafiles\benefits.dta

obs: 1,848

vars: 18

15 Mar 2009 11:25

size: 155,232 (99.9% of memory free)

variable name	storage type	display format	value label	variable label
distid	float	%9.0g		district identifier
schid	int	%9.0g		school identifier
lunch	float	%9.0g		percent eligible, free lunch
enroll	int	%9.0g		school enrollment
staff	float	%9.0g		staff per 1000 students
exppp	int	%9.0g		expenditures per pupil
avgsal	float	%9.0g		average teacher salary, \$
avgben	int	%9.0g		average teacher non-salary benefits, \$
math4	float	%9.0g		percent passing 4th grade math test
story4	float	%9.0g		percent passing 4th grade reading test
bs	float	%9.0g		avgben/avgsal
lavgsal	float	%9.0g		log(avgsal)
lenroll	float	%9.0g		log(enroll)
lstaff	float	%9.0g		log(staff)

Sorted by: distid schid

```
. reg lavgsal bs lstaff lenroll lunch
```

Source	SS	df	MS			
Model	48.3485452	4	12.0871363	Number of obs =	1848	
Residual	51.8328336	1843	.028124164	F(4, 1843) =	429.78	
				Prob > F =	0.0000	
				R-squared =	0.4826	
				Adj R-squared =	0.4815	
Total	100.181379	1847	.054240054	Root MSE =	.1677	

lavgsal	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
bs	-.1774396	.1219691	-1.45	0.146	-.4166518	.0617725
lstaff	-.6907025	.0184598	-37.42	0.000	-.7269068	-.6544981
lenroll	-.0292406	.0084997	-3.44	0.001	-.0459107	-.0125705
lunch	-.0008471	.0001625	-5.21	0.000	-.0011658	-.0005284
_cons	13.72361	.1121095	122.41	0.000	13.50374	13.94349

```
. reg lavgsal bs lstaff lenroll lunch, cluster(distid)
```

(Std. Err. adjusted for 537 clusters in distid)

lavgsal	Coef.	Robust Std. Err.	t	P> t	[95% Conf. Interval]	
bs	-.1774396	.2596214	-0.68	0.495	-.6874398	.3325605
lstaff	-.6907025	.0352962	-19.57	0.000	-.7600383	-.6213666
lenroll	-.0292406	.0257414	-1.14	0.256	-.079807	.0213258
lunch	-.0008471	.0005709	-1.48	0.138	-.0019686	.0002744
_cons	13.72361	.2562909	53.55	0.000	13.22016	14.22707

```
. reg lavgsal bs, cluster(distid)
```

Linear regression

Number of obs = 1848
F(1, 536) = 2.36
Prob > F = 0.1251
R-squared = 0.0049
Root MSE = .23238

(Std. Err. adjusted for 537 clusters in distid)

lavgsal	Coef.	Robust Std. Err.	t	P> t	[95% Conf. Interval]	
bs	-.5034597	.3277449	-1.54	0.125	-1.147282	.1403623
_cons	10.64757	.1056538	100.78	0.000	10.44003	10.85512

```
. xtreg lavgsal bs lstaff lenroll lunch, re
```

```
Random-effects GLS regression           Number of obs   =       1848
Group variable: distid                 Number of groups =        537
```

```
R-sq:  within = 0.5453                 Obs per group:  min =         1
        between = 0.3852                                     avg  =         3.4
        overall = 0.4671                                     max  =        162
```

```
Random effects u_i ~Gaussian           Wald chi2(4)    =    1890.56
corr(u_i, X) = 0 (assumed)             Prob > chi2    =         0.0000
```

lavgsal	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
bs	-.3812698	.1118678	-3.41	0.001	-.6005267	-.162013
lstaff	-.6174177	.0153587	-40.20	0.000	-.6475202	-.5873151
lenroll	-.0249189	.0075532	-3.30	0.001	-.0397228	-.0101149
lunch	.0002995	.0001794	1.67	0.095	-.0000521	.0006511
_cons	13.36682	.0975734	136.99	0.000	13.17558	13.55806
sigma_u	.12627558					
sigma_e	.09996638					
rho	.61473634	(fraction of variance due to u_i)				

```
. xtreg lavgsal bs lstaff lenroll lunch, re cluster(distid)
```

```
Random-effects GLS regression           Number of obs   =       1848
Group variable: distid                 Number of groups =        537
```

```
R-sq:  within = 0.5453                 Obs per group: min =         1
        between = 0.3852                 avg =         3.4
        overall = 0.4671                 max =        162
```

```
Random effects u_i ~Gaussian           Wald chi2(4)    =       316.91
corr(u_i, X) = 0 (assumed)             Prob > chi2     =        0.0000
```

(Std. Err. adjusted for 537 clusters in distid)

lavgsal	Coef.	Robust Std. Err.	z	P> z	[95% Conf. Interval]	
bs	-.3812698	.1504893	-2.53	0.011	-.6762235	-.0863162
lstaff	-.6174177	.0363789	-16.97	0.000	-.688719	-.5461163
lenroll	-.0249189	.0115371	-2.16	0.031	-.0475312	-.0023065
lunch	.0002995	.0001963	1.53	0.127	-.0000852	.0006841
_cons	13.36682	.1968713	67.90	0.000	12.98096	13.75268
sigma_u	.12627558					
sigma_e	.09996638					
rho	.61473634	(fraction of variance due to u_i)				


```
. xtreg lavgsal bs lstaff lenroll lunch, fe
```

```
Fixed-effects (within) regression      Number of obs      =      1848
Group variable: distid                 Number of groups   =       537

R-sq:  within  = 0.5486                 Obs per group:  min =         1
      between  = 0.3544                                     avg  =         3.4
      overall  = 0.4567                                     max  =        162

corr(u_i, Xb) = 0.1433                   F(4,1307)          =    397.05
                                           Prob > F           =     0.0000
```

lavgsal	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
bs	-.4948449	.133039	-3.72	0.000	-.7558382	-.2338515
lstaff	-.6218901	.0167565	-37.11	0.000	-.6547627	-.5890175
lenroll	-.0515063	.0094004	-5.48	0.000	-.0699478	-.0330648
lunch	.0005138	.0002088	2.46	0.014	.0001042	.0009234
_cons	13.61783	.1133406	120.15	0.000	13.39548	13.84018
sigma_u	.15491886					
sigma_e	.09996638					
rho	.70602068	(fraction of variance due to u_i)				
F test that all u_i=0:			F(536, 1307) =	7.24	Prob > F = 0.0000	

```
. xtreg lavgsal bs lstaff lenroll lunch, fe cluster(distid)
```

```
Fixed-effects (within) regression      Number of obs      =      1848
Group variable: distid                 Number of groups   =       537

R-sq:  within  = 0.5486                 Obs per group: min =         1
       between = 0.3544                 avg   =         3.4
       overall  = 0.4567                 max   =        162

                                         F(4,536)          =       57.84
corr(u_i, Xb) = 0.1433                  Prob > F           =       0.0000
```

(Std. Err. adjusted for 537 clusters in distid)

lavgsal	Coef.	Robust Std. Err.	t	P> t	[95% Conf. Interval]	
bs	-.4948449	.1937316	-2.55	0.011	-.8754112	-.1142785
lstaff	-.6218901	.0431812	-14.40	0.000	-.7067152	-.5370649
lenroll	-.0515063	.0130887	-3.94	0.000	-.0772178	-.0257948
lunch	.0005138	.0002127	2.42	0.016	.0000959	.0009317
_cons	13.61783	.2413169	56.43	0.000	13.14379	14.09187
sigma_u	.15491886					
sigma_e	.09996638					
rho	.70602068	(fraction of variance due to u_i)				

```
. xtreg lavgsal bs lstaff lenroll lunch, re cluster(distid) theta
```

```
Random-effects GLS regression           Number of obs   =       1848
Group variable: distid                  Number of groups =        537
```

```
Random effects u_i ~Gaussian           Wald chi2(4)     =       316.91
corr(u_i, X) = 0 (assumed)             Prob > chi2      =        0.0000
```

```
-----+----- theta -----+-----
min      5%      median      95%      max
0.3793   0.3793   0.3793   0.7572   0.9379
```

(Std. Err. adjusted for 537 clusters in distid)

-----+-----						
lavgsal	Coef.	Robust Std. Err.	z	P> z	[95% Conf. Interval]	
-----+-----						
bs	-.3812698	.1504893	-2.53	0.011	-.6762235	-.0863162
lstaff	-.6174177	.0363789	-16.97	0.000	-.688719	-.5461163
lenroll	-.0249189	.0115371	-2.16	0.031	-.0475312	-.0023065
lunch	.0002995	.0001963	1.53	0.127	-.0000852	.0006841
_cons	13.36682	.1968713	67.90	0.000	12.98096	13.75268
-----+-----						
sigma_u	.12627558					
sigma_e	.09996638					
rho	.61473634	(fraction of variance due to u_i)				
-----+-----						

```
. * Create within-district means of all covariates.  
  
. egen bsbar = mean(bs), by(distid)  
. egen lstaffbar = mean(lstaff), by(distid)  
. egen lenrollbar = mean(lenroll), by(distid)  
. egen lunchbar = mean(lunch), by(distid)
```

```
. xtreg lavgsal bs lstaff lenroll lunch bsbar lstaffbar lenrollbar lunchbar,
    re cluster(distid)
```

```
Random-effects GLS regression           Number of obs   =       1848
Group variable: distid                  Number of groups =        537
```

(Std. Err. adjusted for 537 clusters in distid)

lavgsal	Coef.	Robust Std. Err.	z	P> z	[95% Conf. Interval]	
bs	-.4948449	.1939422	-2.55	0.011	-.8749646	-.1147252
lstaff	-.6218901	.0432281	-14.39	0.000	-.7066157	-.5371645
lenroll	-.0515063	.013103	-3.93	0.000	-.0771876	-.025825
lunch	.0005138	.000213	2.41	0.016	.0000964	.0009312
bsbar	.2998553	.3031961	0.99	0.323	-.2943981	.8941088
lstaffbar	-.0255493	.0651932	-0.39	0.695	-.1533256	.1022269
lenrollbar	.0657285	.020655	3.18	0.001	.0252455	.1062116
lunchbar	-.0007259	.0004378	-1.66	0.097	-.0015839	.0001322
_cons	13.22003	.2556139	51.72	0.000	12.71904	13.72103
sigma_u	.12627558					
sigma_e	.09996638					
rho	.61473633	(fraction of variance due to u_i)				

```
. test bsbar lstaffbar lenrollbar lunchbar
```

```
( 1) bsbar = 0  
( 2) lstaffbar = 0  
( 3) lenrollbar = 0  
( 4) lunchbar = 0
```

```
      chi2( 4) =    20.70  
Prob > chi2 =    0.0004
```

2. Cluster-Robust Inference with Large Group Sizes

- What if one applies robust inference when the fixed M_g , $G \rightarrow \infty$ asymptotic analysis is not realistic? If the M_g are “large” along with G , valid inference is still possible.
- Hansen (2007, Theorem 2, *Journal of Econometrics*) shows that with G and M_g both getting large the usual inference based on the robust “sandwich” estimator is valid with arbitrary correlation among the errors, v_{gm} , within each group. (Independence across groups is maintained.)

- For example, if we have a sample of $G = 100$ schools and roughly $M_g = 100$ students per school cluster-robust inference for pooled OLS should produce inference of roughly the correct size.

- Unfortunately, in the presence of cluster effects with a small number of groups (G) and large group sizes (M_g), cluster-robust inference with pooled OLS falls outside Hansen's theoretical findings. We should not expect good properties of the cluster-robust inference with small groups and large group sizes.

- Example: Suppose $G = 10$ hospitals have been sampled with several hundred patients per hospital. If the explanatory variable of interest varies only at the hospital level, tempting to use pooled OLS with cluster-robust inference. But we have no theoretical justification for doing so, and reasons to expect it will not work well.

- If the explanatory variables of interest vary within group, FE is attractive. First, allows c_g to be arbitrarily correlated with the \mathbf{z}_{gm} . Second, with large M_g , can treat the c_g as parameters to estimate – because we can estimate them precisely – and then assume that the observations are independent across m (as well as g). This means that the usual inference is valid, perhaps with adjustment for heteroskedasticity.

- For panel data applications, Hansen's (2007) results, particularly Theorem 3, imply that cluster-robust inference for the fixed effects estimator should work well when the cross section (N) and time series (T) dimensions are similar and not too small. If full time effects are allowed in addition to unit-specific fixed effects – as they often should – then the asymptotics must be with N and T both getting large.

- Any serial dependence in the idiosyncratic errors is assumed to be weakly dependent. Simulations in Bertrand, Duflo, and Mullainathan (2004) and Hansen (2007) verify that the robust cluster-robust variance matrix works well when N and T are about 50 and the idiosyncratic errors follow a stable AR(1) model.

3. Cluster Samples with Unit-Specific Panel Data

- Often, cluster samples come with a time component, so that there are two potential sources of correlation across observations: across time within the same individual and across individuals within the same group.
- Assume here that there is a natural nesting. Each unit belongs to a cluster and the cluster identification does not change over time.
- For example, we might have annual panel data at the firm level, and each firm belongs to the same industry (cluster) for all years. Or, we have panel data for schools that each belong to a district.

- Special case of **hierarchical linear model (HLM)** setup or **mixed models** or **multilevel models**.
- Now we have three data subscripts on at least some variables that we observe. For example, the response variable is y_{gmt} , where g indexes the group or cluster, m is the unit within the group, and t is the time index.
- Assume we have a balanced panel with the time periods running from $t = 1, \dots, T$. (Unbalanced case not difficult, assuming exogenous selection.) Within cluster g there are M_g units, and we have sampled G clusters. (In the HLM literature, g is usually called the *first level* and m the *second level*.)

- We assume that we have many groups, G , and relatively few members of the group. Asymptotics: fixed M_g and T fixed with G getting large. For example, if we can sample, say, several hundred school districts, with a few to maybe a few dozen schools per district, over a handful of years, then we have a data set that can be analyzed in the current framework.

- A standard linear model with constant slopes can be written, for $t = 1, \dots, T$, $m = 1, \dots, M_g$, and a random draw g from the population of clusters as

$$y_{gmt} = \eta_t + \mathbf{w}_g \boldsymbol{\alpha} + \mathbf{x}_{gm} \boldsymbol{\beta} + \mathbf{z}_{gmt} \boldsymbol{\delta} + h_g + c_{gm} + u_{gmt},$$

where, say, h_g is the industry or district effect, c_{gm} is the firm effect or school effect (firm or school m in industry or district g), and u_{gmt} is the idiosyncratic effect. In other words, the composite error is

$$v_{gmt} = h_g + c_{gm} + u_{gmt}.$$

- Generally, the model can include variables that change at any level.
- Some elements of \mathbf{z}_{gmt} might change only across g and t , and not by unit. This is an important special case for policy analysis where the policy applies at the group level but changes over time.
- With the presence of \mathbf{w}_g , or variables that change across g and t , need to recognize h_g .

- If assume the error v_{gmt} is uncorrelated with $(\mathbf{w}_g, \mathbf{x}_{gm}, \mathbf{z}_{gmt})$, pooled OLS is simple and attractive. Consistent as $G \rightarrow \infty$ for any cluster or serial correlation pattern.
- The most general inference for pooled OLS – still maintaining independence across clusters – is to allow any kind of serial correlation across units or time, or both, within a cluster.

- In Stata:

```
reg y w1 ... wJ x1 ... xK z1 ... zL,  
    cluster(industryid)
```

- Compare with inference robust only to serial correlation:

```
reg y w1 ... wJ x1 ... xK z1 ... zL,  
    cluster(firmid)
```

- In the context of cluster sampling with panel data, the latter is no longer “fully robust” because it ignores possible within-cluster correlation.

- Can apply a generalized least squares analysis that makes assumptions about the components of the composite error. Typically, assume components are pairwise uncorrelated, the c_{gm} are uncorrelated within cluster (with common variance), and the u_{gmt} are uncorrelated within cluster and across time (with common variance).
- Resulting feasible GLS estimator is an extension of the usual random effects estimator for panel data.
- Because of the large- G setting, the estimator is consistent and asymptotically normal whether or not the actual variance structure we use in estimation is the proper one.

- To guard against heteroskedasticity in any of the errors and serial correlation in the $\{u_{gmt}\}$, one should use fully robust inference that does not rely on the form of the unconditional variance matrix (which may also differ from the conditional variance matrix).
- Simpler strategy: apply random effects at the individual level, effectively ignoring the clusters *in estimation*. In other words, treat the data as a standard panel data set in estimation and apply usual RE. To account for the cluster sampling in inference, one computes a fully robust variance matrix estimator for the usual random effects estimator.

- In Stata:

```
xtset firmid year
```

```
xtreg y w1 ... wJ x1 ... xK z1 ... zL, re  
      cluster(industryid)
```

- Again, compare with inference robust only to neglected serial correlation:

```
xtreg y w1 ... wJ x1 ... xK z1 ... zL, re  
      cluster(firmid)
```

- Formal analysis. Write the equation for each cluster as

$$\mathbf{y}_g = \mathbf{R}_g \boldsymbol{\theta} + \mathbf{v}_g$$

where a row of \mathbf{R}_g is $(1, d_2, \dots, d_T, \mathbf{w}_g, \mathbf{x}_{gm}, \mathbf{z}_{gmt})$ (which includes a full set of period dummies) and $\boldsymbol{\theta}$ is the vector of all regression parameters. For cluster g , \mathbf{y}_g contains $M_g T$ elements (T periods for each unit m).

- In particular,

$$\mathbf{y}_g = \begin{pmatrix} \mathbf{y}_{g1} \\ \mathbf{y}_{g2} \\ \vdots \\ \mathbf{y}_{g,M_g} \end{pmatrix}, \quad \mathbf{y}_{gm} = \begin{pmatrix} y_{gm1} \\ y_{gm2} \\ \vdots \\ y_{gmT} \end{pmatrix}$$

so that each \mathbf{y}_{gm} is $T \times 1$; \mathbf{v}_g has an identical structure. Now, we can obtain $\mathbf{\Omega}_g = \text{Var}(\mathbf{v}_g)$ under various assumptions and apply feasible GLS.

- RE at the unit level is obtained by choosing $\mathbf{\Omega}_g = \mathbf{I}_{M_g} \otimes \mathbf{\Lambda}$, where $\mathbf{\Lambda}$ is the $T \times T$ matrix with the RE structure. If there is within-cluster correlation, this is not the correct form of $Var(\mathbf{v}_g)$, and that is why robust inference is generally needed after RE estimation.

- For the case that $v_{gmt} = h_g + c_{gm} + u_{gmt}$ where the terms have variances σ_h^2 , σ_c^2 , and σ_u^2 , respectively, they are pairwise uncorrelated, c_{gm} and c_{gr} are uncorrelated for $r \neq m$, and $\{u_{gmt} : t = 1, \dots, T\}$ is serially uncorrelated, we can obtain $\mathbf{\Omega}_g$ as follows:

$$\text{Var}(\mathbf{v}_{gm}) = (\sigma_h^2 + \sigma_c^2)\mathbf{j}_T\mathbf{j}'_T + \sigma_u^2\mathbf{I}_T$$

$$\text{Cov}(\mathbf{v}_{gm}, \mathbf{v}_{gr}) = \sigma_h^2\mathbf{j}_T\mathbf{j}'_T, r \neq m$$

$$\mathbf{\Omega}_g = \begin{pmatrix} (\sigma_h^2 + \sigma_c^2)\mathbf{j}_T\mathbf{j}'_T + \sigma_u^2\mathbf{I}_T & \cdots & \sigma_h^2\mathbf{j}_T\mathbf{j}'_T \\ \vdots & \ddots & \vdots \\ \sigma_h^2\mathbf{j}_T\mathbf{j}'_T & \cdots & (\sigma_h^2 + \sigma_c^2)\mathbf{j}_T\mathbf{j}'_T + \sigma_u^2\mathbf{I}_T \end{pmatrix}$$

- The robust asymptotic variance of $\hat{\boldsymbol{\theta}}$ is estimated as

$$\widehat{Avar}(\hat{\boldsymbol{\theta}}) = \left(\sum_{g=1}^G \mathbf{R}'_g \hat{\boldsymbol{\Omega}}_g^{-1} \mathbf{R}_g \right)^{-1} \left(\sum_{g=1}^G \mathbf{R}'_g \hat{\boldsymbol{\Omega}}_g^{-1} \hat{\mathbf{v}}_g \hat{\mathbf{v}}'_g \hat{\boldsymbol{\Omega}}_g^{-1} \mathbf{R}_g \right)^{-1} \cdot \left(\sum_{g=1}^G \mathbf{R}'_g \hat{\boldsymbol{\Omega}}_g^{-1} \mathbf{R}_g \right)^{-1},$$

where $\hat{\mathbf{v}}_g = \mathbf{y}_g - \mathbf{R}_g \hat{\boldsymbol{\theta}}$.

- Unfortunately, routines intended for estimating HLMs (or mixed models) assume that the structure imposed on $\mathbf{\Omega}_g$ is correct, and that $Var(\mathbf{v}_g|\mathbf{R}_g) = Var(\mathbf{v}_g)$. The resulting inference could be misleading, especially if serial correlation in $\{u_{gmt}\}$ is not allowed.
- In Stata, the command is `xtmixed`.

- Because of the nested data structure, we have available different versions of fixed effects estimators. Subtracting cluster averages from all observations within a cluster eliminates h_g ; when $\mathbf{w}_{gt} = \mathbf{w}_g$ for all t , \mathbf{w}_g is also eliminated. But the unit-specific effects, c_{mg} , are still part of the error term. If we are mainly interested in δ , the coefficients on the time-varying variables \mathbf{z}_{gmt} , then removing c_{gm} (along with h_g) is attractive. In other words, use a standard fixed effects analysis at the individual level.

- If the units are allowed to change groups over time – such as children changing schools – then we would replace h_g with h_{gt} , and then subtracting off individual-specific means would not remove the time-varying cluster effects.

- Even if we use unit “fixed effects” – that is, we demean the data at the unit level – we might still use inference robust to clustering at the aggregate level. Suppose the model is

$$\begin{aligned}y_{gmt} &= \eta_t + \mathbf{w}_g \boldsymbol{\alpha} + \mathbf{x}_{gm} \boldsymbol{\beta} + \mathbf{z}_{gmt} \mathbf{d}_{mg} + h_g + c_{mg} + u_{gmt} \\ &= \eta_t + \mathbf{w}_{gt} \boldsymbol{\alpha} + \mathbf{x}_{gm} \boldsymbol{\beta} + \mathbf{z}_{gmt} \boldsymbol{\delta} + h_g + c_{mg} + u_{gmt} + \mathbf{z}_{gmt} \mathbf{e}_{gm},\end{aligned}$$

where $\mathbf{d}_{gm} = \boldsymbol{\delta} + \mathbf{e}_{gm}$ is a set of unit-specific intercepts on the individual, time-varying covariates \mathbf{z}_{gmt} .

- The time-demeaned equation within individual m in cluster g is

$$y_{gmt} - \bar{y}_{gm} = \zeta_t + (\mathbf{z}_{gmt} - \bar{\mathbf{z}}_{gm})\boldsymbol{\delta} + (u_{gmt} - \bar{u}_{gm}) + (\mathbf{z}_{gmt} - \bar{\mathbf{z}}_{gm})\mathbf{e}_{gm}.$$

- FE is still consistent if $E(\mathbf{d}_{mg} | \mathbf{z}_{gmt} - \bar{\mathbf{z}}_{gm}) = E(\mathbf{d}_{mg})$, $m = 1, \dots, M_g$, $t = 1, \dots, T$, and all g , and so cluster-robust inference, which is automatically robust to serial correlation and heteroskedsticity, makes perfectly good sense.

● Example: Effects of Funding on Student Performance

```
. use meap94_98
```

```
. des
```

Contains data from meap94_98.dta

```
obs:      7,150
vars:      26                               13 Mar 2009 11:30
size:      893,750 (99.8% of memory free)
```

```
-----
```

variable name	storage type	display format	value label	variable label
distid	float	%9.0g		district identifier
schid	int	%9.0g		school identifier
lunch	float	%9.0g		% eligible for free lunch
enrol	int	%9.0g		number of students
exppp	int	%9.0g		expenditure per pupil
math4	float	%9.0g		% satisfactory, 4th grade math test
year	int	%9.0g		1992=school yr 1991-2
cpi	float	%9.0g		consumer price index
rexppp	float	%9.0g		(exppp/cpi)*1.695: 1997 \$
lrexpp	float	%9.0g		log(rexpp)
lenrol	float	%9.0g		log(enrol)
avgrexp	float	%9.0g		(rexppp + rexppp_1)/2
lavgrexp	float	%9.0g		log(avgrexp)
tobs	byte	%9.0g		number of time periods

```
-----
```

```
Sorted by:  schid  year
```

```
. * egen tobs = sum(1), by(schid)
```

```
. tab tobs if y98
```

number of time periods	Freq.	Percent	Cum.
3	487	29.28	29.28
4	254	15.27	44.56
5	922	55.44	100.00
Total	1,663	100.00	

```
. xtreg math4 lavgrexp lunch lenrol y95-y98, fe
```

```
Fixed-effects (within) regression      Number of obs      =      7150
Group variable: schid                  Number of groups   =      1683

R-sq:  within  = 0.3602                Obs per group: min =      3
       between = 0.0292                avg   =      4.2
       overall  = 0.1514                max   =      5
```

math4	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
lavgrexp	6.288376	2.098685	3.00	0.003	2.174117	10.40264
lunch	-.0215072	.0312185	-0.69	0.491	-.082708	.0396935
lenrol	-2.038461	1.791604	-1.14	0.255	-5.550718	1.473797
y95	11.6192	.5545233	20.95	0.000	10.53212	12.70629
y96	13.05561	.6630948	19.69	0.000	11.75568	14.35554
y97	10.14771	.7024067	14.45	0.000	8.770713	11.52471
y98	23.41404	.7187237	32.58	0.000	22.00506	24.82303
_cons	11.84422	22.81097	0.52	0.604	-32.87436	56.5628
sigma_u	15.84958					
sigma_e	11.325028					
rho	.66200804	(fraction of variance due to u_i)				

```
F test that all u_i=0:      F(1682, 5460) =      4.82      Prob > F = 0.0000
```

```
. xtreg math4 lavgrexp lunch lenrol y95-y98, fe cluster(schid)
```

```
Fixed-effects (within) regression      Number of obs      =      7150
Group variable: schid                  Number of groups   =      1683
```

(Std. Err. adjusted for 1683 clusters in schid)

math4	Coef.	Robust Std. Err.	t	P> t	[95% Conf. Interval]	
lavgrexp	6.288376	2.431317	2.59	0.010	1.519651	11.0571
lunch	-.0215072	.0390732	-0.55	0.582	-.0981445	.05513
lenrol	-2.038461	1.789094	-1.14	0.255	-5.547545	1.470623
y95	11.6192	.5358469	21.68	0.000	10.56821	12.6702
y96	13.05561	.6910815	18.89	0.000	11.70014	14.41108
y97	10.14771	.7326314	13.85	0.000	8.710745	11.58468
y98	23.41404	.7669553	30.53	0.000	21.90975	24.91833
_cons	11.84422	25.16643	0.47	0.638	-37.51659	61.20503
sigma_u	15.84958					
sigma_e	11.325028					
rho	.66200804	(fraction of variance due to u_i)				

```
. xtreg math4 lavgrexp lunch lenrol y95-y98, fe cluster(distid)
```

(Std. Err. adjusted for 467 clusters in distid)

math4	Coef.	Robust Std. Err.	t	P> t	[95% Conf. Interval]	
lavgrexp	6.288376	3.132334	2.01	0.045	.1331271	12.44363
lunch	-.0215072	.0399206	-0.54	0.590	-.0999539	.0569395
lenrol	-2.038461	2.098607	-0.97	0.332	-6.162365	2.085443
y95	11.6192	.7210398	16.11	0.000	10.20231	13.0361
y96	13.05561	.9326851	14.00	0.000	11.22282	14.8884
y97	10.14771	.9576417	10.60	0.000	8.26588	12.02954
y98	23.41404	1.027313	22.79	0.000	21.3953	25.43278
_cons	11.84422	32.68429	0.36	0.717	-52.38262	76.07107
sigma_u	15.84958					
sigma_e	11.325028					
rho	.66200804	(fraction of variance due to u_i)				

4. Clustering and Stratification

- Survey data often characterized by clustering and VP sampling.

Suppose that g represents the primary sampling unit (say, city) and individuals or families (indexed by m) are sampled within each PSU with probability p_{gm} . If $\hat{\beta}$ is the pooled OLS estimator across PSUs and individuals, its variance is estimated as

$$\begin{aligned}
& \left(\sum_{g=1}^G \sum_{m=1}^{M_g} \mathbf{x}'_{gm} \mathbf{x}_{gm} / p_{gm} \right)^{-1} \\
& \cdot \left[\sum_{g=1}^G \sum_{m=1}^{M_g} \sum_{r=1}^{M_g} \hat{u}_{gm} \hat{u}_{gr} \mathbf{x}'_{gm} \mathbf{x}_{gr} / (p_{gm} p_{gr}) \right] \\
& \cdot \left(\sum_{g=1}^G \sum_{m=1}^{M_g} \mathbf{x}'_{gm} \mathbf{x}_{gm} / p_{gm} \right)^{-1} .
\end{aligned}$$

If the probabilities are estimated using retention frequencies, estimate is conservative, as before.

- Multi-stage sampling schemes introduce even more complications.

Let there be S strata (e.g., states in the U.S.), exhaustive and mutually exclusive. Within stratum s , there are C_s clusters (e.g., neighborhoods).

- Large-sample approximations: the number of clusters sampled, N_s , gets large. This allows for arbitrary correlation (say, across households) within cluster.

- Within stratum s and cluster c , let there be M_{sc} total units (household or individuals). Therefore, the total number of units in the population is

$$M = \sum_{s=1}^S \sum_{c=1}^{C_s} M_{sc}.$$

- Let z be a variable whose mean we want to estimate. List all population values as $\{z_{scm}^o : m = 1, \dots, M_{sc}, c = 1, \dots, C_s, s = 1, \dots, S\}$, so the population mean is

$$\mu = M^{-1} \sum_{s=1}^S \sum_{c=1}^{C_s} \sum_{m=1}^{M_{sc}} z_{scm}^o.$$

Define the total in the population as

$$\tau = \sum_{s=1}^S \sum_{c=1}^{C_s} \sum_{m=1}^{M_{sc}} z_{scm}^o = M\mu.$$

Totals within each cluster and then stratum are, respectively,

$$\tau_{sc} = \sum_{m=1}^{M_{sc}} z_{scm}^o$$

$$\tau_s = \sum_{c=1}^{C_s} \tau_{sc}$$

- Sampling scheme:

(i) For each stratum s , randomly draw N_s clusters, with replacement.

(Fine for “large” C_s .)

(ii) For each cluster c drawn in step (i), randomly sample K_{sc} households with replacement.

- For each pair (s, c) , define

$$\hat{\mu}_{sc} = K_{sc}^{-1} \sum_{m=1}^{K_{sc}} z_{scm}.$$

Because this is a random sample within (s, c) ,

$$E(\hat{\mu}_{sc}) = \mu_{sc} = M_{sc}^{-1} \sum_{m=1}^{M_{sc}} z_{scm}^o.$$

- To continue up to the cluster level we need to estimate the total, $\tau_{sc} = M_{sc}\mu_{sc}$. So, $\hat{\tau}_{sc} = M_{sc}\hat{\mu}_{sc}$ is an unbiased estimator of τ_{sc} for all $\{(s, c) : c = 1, \dots, C_s, s = 1, \dots, S\}$. (We can think of computing this estimate even if we eventually do not use some clusters.)

- Next, consider randomly drawing N_s clusters from stratum s . Can show that an unbiased estimator of the total τ_s for stratum s is

$$C_s \cdot N_s^{-1} \sum_{c=1}^{N_s} \hat{\tau}_{sc}.$$

- Finally, the total in the population is estimated as

$$\sum_{s=1}^S \left(C_s \cdot N_s^{-1} \sum_{c=1}^{N_s} \hat{\tau}_{sc} \right) \equiv \sum_{s=1}^S \sum_{c=1}^{N_s} \sum_{m=1}^{K_{sc}} \omega_{sc} Z_{scm}$$

where the weight for stratum-cluster pair (s, c) is

$$\omega_{sc} \equiv \frac{C_s}{N_s} \cdot \frac{M_{sc}}{K_{sc}}.$$

- Note how $\omega_{sc} = (C_s/N_s)(M_{sc}/K_{sc})$ accounts for under- or over-sampled clusters within strata and under- or over-sampled units within clusters.
- Appears in the literature on complex survey sampling, sometimes without M_{sc}/K_{sc} when each cluster is sampled as a complete unit, and so $M_{sc}/K_{sc} = 1$.
- To estimate the mean μ , just divide by M , the total number of units sampled.

$$\hat{\mu} = M^{-1} \left(\sum_{s=1}^S \sum_{c=1}^{N_s} \sum_{m=1}^{K_{sc}} \omega_{sc} z_{scm} \right).$$

- To study regression, specify the problem as

$$\min_{\boldsymbol{\beta}} \sum_{s=1}^S \sum_{c=1}^{N_s} \sum_{m=1}^{K_{sc}} \omega_{sc} (y_{scm} - \mathbf{x}_{scm} \boldsymbol{\beta})^2.$$

The asymptotic variance combines clustering with weighting to account for the multi-stage sampling. Following Bhattacharya (2005, Journal of Econometrics), an appropriate asymptotic variance estimate has a sandwich form,

$$\left(\sum_{s=1}^S \sum_{c=1}^{N_s} \sum_{m=1}^{K_{sc}} \omega_{sc} \mathbf{x}'_{scm} \mathbf{x}_{scm} \right)^{-1} \hat{\mathbf{B}} \left(\sum_{s=1}^S \sum_{c=1}^{N_s} \sum_{m=1}^{K_{sc}} \omega_{sc} \mathbf{x}'_{scm} \mathbf{x}_{scm} \right)^{-1}.$$

- $\hat{\mathbf{B}}$ is somewhat complicated:

$$\begin{aligned}
\hat{\mathbf{B}} = & \sum_{s=1}^S \sum_{c=1}^{N_s} \sum_{m=1}^{K_{sc}} \omega_{sc}^2 \hat{u}_{scm}^2 \mathbf{x}'_{scm} \mathbf{x}_{scm} \\
& + \sum_{s=1}^S \sum_{c=1}^{N_s} \sum_{m=1}^{K_{sc}} \sum_{r \neq m}^{K_{sc}} \omega_{sc}^2 \hat{u}_{scm} \hat{u}_{scr} \mathbf{x}'_{scm} \mathbf{x}_{scr} \\
& - \sum_{s=1}^S N_s^{-1} \left(\sum_{c=1}^{N_s} \sum_{m=1}^{K_{sc}} \omega_{sc} \mathbf{x}'_{scm} \hat{u}_{scm} \right) \left(\sum_{c=1}^{N_s} \sum_{m=1}^{K_{sc}} \omega_{sc} \mathbf{x}'_{scm} \hat{u}_{scm} \right)'
\end{aligned}$$

- The first part of $\hat{\mathbf{B}}$ is obtained using the White “heteroskedasticity”-robust form. The second piece accounts for the clustering. The third piece reduces the variance by accounting for the nonzero means of the “score” within strata.

- Suppose that the population is stratified by region, taking on values 1 through 8, and the primary sampling unit is zip code. Within each zip code we obtain a sample of families, possibly using VP sampling.

- Stata command:

```
svyset zipcode [pweight = sampwght],  
strata(region)
```

- Now we can use a set of econometric commands. For example,

```
svy: reg y x1 ... xK
```

```

. use http://www.stata-press.com/data/r10/nhanes2f

. svyset psuid [pweight = finalwgt], strata(stratid)
pweight: finalwgt
VCE: linearized
Single unit: missing
Strata 1: stratid
SU 1: psuid
FPC 1: <zero>

```

```

. tab health

```

1=excellent , . . . , 5=poor	Freq.	Percent	Cum.
poor	729	7.05	7.05
fair	1,670	16.16	23.21
average	2,938	28.43	51.64
good	2,591	25.07	76.71
excellent	2,407	23.29	100.00
Total	10,335	100.00	

```

. sum lead

```

Variable	Obs	Mean	Std. Dev.	Min	Max
lead	4942	14.32032	6.167695	2	80

```
. svy: oprobit health lead female black age weight
(running oprobit on estimation sample)
```

Survey: Ordered probit regression

```
Number of strata   =          31      Number of obs       =          4940
Number of PSUs     =          62      Population size     =       56316764
                                          Design df          =           31
                                          F( 5, 27)         =          78.49
                                          Prob > F          =          0.0000
```

health	Coef.	Linearized Std. Err.	t	P> t	[95% Conf. Interval]	
lead	-.0059646	.0045114	-1.32	0.196	-.0151656	.0032364
female	-.1529889	.057348	-2.67	0.012	-.2699508	-.036027
black	-.535801	.0622171	-8.61	0.000	-.6626937	-.4089084
age	-.0236837	.0011995	-19.75	0.000	-.02613	-.0212373
weight	-.0035402	.0010954	-3.23	0.003	-.0057743	-.0013061
/cut1	-3.278321	.1711369	-19.16	0.000	-3.627357	-2.929285
/cut2	-2.496875	.1571842	-15.89	0.000	-2.817454	-2.176296
/cut3	-1.611873	.1511986	-10.66	0.000	-1.920244	-1.303501
/cut4	-.8415657	.1488381	-5.65	0.000	-1.145123	-.5380083

```
. oprobit health lead female black age weight
```

```
Iteration 0: log likelihood = -7526.7772  
Iteration 1: log likelihood = -7133.9477  
Iteration 2: log likelihood = -7133.6805
```

```
Ordered probit regression
```

```
Number of obs = 4940  
LR chi2(5) = 786.19  
Prob > chi2 = 0.0000  
Pseudo R2 = 0.0522
```

```
Log likelihood = -7133.6805
```

health	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
lead	-.0011088	.0026942	-0.41	0.681	-.0063893	.0041718
female	-.1039273	.0352721	-2.95	0.003	-.1730594	-.0347952
black	-.4942909	.0502051	-9.85	0.000	-.592691	-.3958908
age	-.0237787	.0009147	-26.00	0.000	-.0255715	-.0219859
weight	-.0027245	.0010558	-2.58	0.010	-.0047938	-.0006551
/cut1	-3.072779	.1087758			-3.285975	-2.859582
/cut2	-2.249324	.1057841			-2.456657	-2.041991
/cut3	-1.396732	.1038044			-1.600185	-1.19328
/cut4	-.6615336	.1028773			-.8631693	-.4598978

5. Two-Way Clustering

- Recent interest in two-way clustering – usually across time within a firm and across firms within a given time period.
- Often the underlying model is set up as follows:

$$y_{it} = \mathbf{x}_{it}\boldsymbol{\beta} + g_t + c_i + u_{it}$$

where g_t and c_i are both viewed as random (as is u_{it} , of course).

- The presence of c_i , as usual, induces correlation across time. But $\{u_{it}\}$ can also be serially correlated across across time.

- If the firm effects c_i and idiosyncratic errors $\{u_{it}\}$ are uncorrelated across i – as in many data generating mechanisms – then all cross-sectional correlation is due to the presence of g_t , the aggregate time effects.
- Eliminating g_t by the within transformation across firms in the same time period solves the problem. In practice, include time dummies along with firm dummies. Or, include time dummies and use standard FE software, as in

```
xtset firmid year  
  
xi: xtreg y x1 ... xK i.year, fe cluster(firmid)
```

- Could there be left over cross-sectional correlation even if we use time dummies? Yes, for example if

$$y_{it} = \mathbf{x}_{it}\mathbf{b}_t + g_t + c_i + u_{it}$$

but we act as if $\mathbf{b}_t = \boldsymbol{\beta}$. [Then $\mathbf{x}_{it}(\mathbf{b}_t - \boldsymbol{\beta})$ is part of the error term.] But how important is this in practice?

- Peterson (2009, Review of Financial Studies) and Gow, Ormazabal, and Taylor (2010, Accounting Review) study various standard errors, including two-way clustering, for simulated and actual data. The two-way clustering appears to work well even for T as small as 10 and $N = 200$.

- But some empirical examples in GOT are misleading. With very large N and relatively small T (15 to 30), they compare two-way clustering with clustering for cross-sectional correlation. The obvious approach is to include year dummies (which they do in some cases) and cluster for serial correlation. Firm effects may or may not be needed for consistency of the parameters, but including them would reduce the time series correlation. GOT do not include them.
- Thompson (2011, Journal of Financial Economics) shows that the two-way clustering (or double clustering) is valid provided N and T are both “large.” The aggregate shocks must dissipate over time.

- In simulations, two-way clustering seems to work well for $N = 50$ and $T = 25$, but there is little justification for clustering for cross-sectional correlation if, say, $N = 1,000$ and $T = 5$.
- In such scenarios, the two-way clustering actually produces standard errors that are too small if there is no time series nor cross-sectional correlation.
- With large N and small T , it seems including time effects and clustering for serial correlation is the only theoretically justified procedure.

● Example: Airfare equation:

. * First, two-way clustering without firm effects.

```
. xi: cluster2 lfare concen ldist ldistsq i.year, fcluster(id) tcluster(year)
i.year          _Iyear_1997-2000      (naturally coded; _Iyear_1997 omitted)
```

Linear regression with 2D clustered SEs

Number of obs = 4596
 F(6, 4589) = 558.39
 Prob > F = 0.0000
 R-squared = 0.4062
 Root MSE = 0.3365

Number of clusters (id) = 1149
 Number of clusters (year) = 4

	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
concen	.3601203	.0560493	6.43	0.000	.2502368	.4700039
ldist	-.9016004	.235178	-3.83	0.000	-1.362662	-.4405384
ldistsq	.1030196	.0174188	5.91	0.000	.0688704	.1371688
_Iyear_1998	.0211244
_Iyear_1999	.0378496
_Iyear_2000	.09987
_cons	6.209258	.7956274	7.80	0.000	4.649445	7.76907

SE clustered by id and year

. * Now cluster only within route to account for the substantial serial
. * correlation. Remember, the route effect is left in the error term.

```
. xi: reg lfare concen ldist ldistsq i.year, cluster(id)
i.year      _Iyear_1997-2000    (naturally coded; _Iyear_1997 omitted)
```

```
Linear regression                               Number of obs =    4596
                                                F(   6,  1148) =   205.63
                                                Prob > F       =    0.0000
                                                R-squared      =    0.4062
                                                Root MSE      =    .33651
```

(Std. Err. adjusted for 1149 clusters in id)

lfare	Coef.	Robust Std. Err.	t	P> t	[95% Conf. Interval]	
concen	.3601203	.058556	6.15	0.000	.2452315	.4750092
ldist	-.9016004	.2719464	-3.32	0.001	-1.435168	-.3680328
ldistsq	.1030196	.0201602	5.11	0.000	.0634647	.1425745
_Iyear_1998	.0211244	.0041474	5.09	0.000	.0129871	.0292617
_Iyear_1999	.0378496	.0051795	7.31	0.000	.0276872	.048012
_Iyear_2000	.09987	.0056469	17.69	0.000	.0887906	.1109493
_cons	6.209258	.9117551	6.81	0.000	4.420364	7.998151

. * These are much closer to the two-way cluster standard errors; in fact,
. * somewhat larger. And these have justification with large N and small T.

. * What if we also use firm FEs (on a reduced sample)?

```
. xi: xtreg lfare concen ldist ldistsq i.year, fe cluster(id)
i.year          _Iyear_1997-2000    (naturally coded; _Iyear_1997 omitted)
```

```
Fixed-effects (within) regression      Number of obs      =      400
Group variable: id                    Number of groups   =      100
```

```
corr(u_i, Xb) = -0.3263                F(4,99)           =      11.52
                                          Prob > F          =      0.0000
```

(Std. Err. adjusted for 100 clusters in id)

lfare	Coef.	Robust Std. Err.	t	P> t	[95% Conf. Interval]	
concen	.5585469	.2097257	2.66	0.009	.1424057	.9746881
ldist	(dropped)					
ldistsq	(dropped)					
_Iyear_1998	-.0043007	.0185779	-0.23	0.817	-.0411632	.0325618
_Iyear_1999	.0324459	.0200249	1.62	0.108	-.0072878	.0721797
_Iyear_2000	.0878409	.0206729	4.25	0.000	.0468213	.1288604
_cons	4.675322	.1408638	33.19	0.000	4.395817	4.954826
sigma_u	.37074456					
sigma_e	.11866722					
rho	.90707061	(fraction of variance due to u_i)				


```

. xi: cluster2 lfare concen ldist ldistsq i.id i.year, fcluster(id)
      tcluster(year)
i.id      _Iid_1-100      (naturally coded; _Iid_1 omitted)
i.year    _Iyear_1997-2000 (naturally coded; _Iyear_1997 omitted)

```

```

Linear regression with 2D clustered SEs
Number of obs = 400
F(103, 296) = 229.75
Prob > F = 0.0000
R-squared = 0.9211
Root MSE = 0.1187
Number of clusters (id) = 100
Number of clusters (year) = 4

```

	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
concen	.5585469	.2756093	2.03	0.044	.0161449	1.100949
ldist	(dropped)					
ldistsq	.0535631	.0007484	71.57	0.000	.0520902	.0550361
_Iid_2	-.2053579	.0484883	-4.24	0.000	-.3007833	-.1099324
_Iid_3	.3550181
.... (output suppressed)						

SE clustered by id and year

```

. * Now the two-way standard error is quite a bit larger. But we have no theory
. * telling us it is valid.

```

- The “cluster2” command is tied to structures such as

$$y_{it} = \mathbf{x}_{it}\boldsymbol{\beta} + g_t + c_i + u_{it}$$

It does not allow situations where, say, shocks to firm h in year $t - 1$ are correlated with shocks to firm $i \neq h$ in year t .

- Thompson (2011) allows for some dependence across time for different units, but must specify the maximum lag.